





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : A61K 31/41, 31/4164, 31/4178, 31/4196, 31/4245, C07D 233/54, 233/56, 233/60, 233/68, 249/12, 254/10, 263/30, 263/34, 403/00</p>	A1	<p>(11) International Publication Number: <b>WO 00/59506</b></p> <p>(43) International Publication Date: 12 October 2000 (12.10.00)</p>
<p>(21) International Application Number: PCT/US00/07417</p> <p>(22) International Filing Date: 20 March 2000 (20.03.00)</p> <p>(30) Priority Data: 60/127,745 5 April 1999 (05.04.99) US</p> <p>(71) Applicant: BRISTOL-MYERS SQUIBB CO. [US/US]; P.O. Box 4000, Princeton, NJ 08543 (US).</p> <p>(72) Inventors: ROBL, Jeffrey, A.; 7 Tulip Drive, Newtown, PA 18940 (US). SULSKY, Richard, B.; 2 Coventry Square, West Trenton, NJ 08623 (US). MAGNIN, David, R.; 40 Cottage Court, Hamilton, NJ 08690 (US).</p> <p>(74) Agents: DAVIS, Stephen, B. et al.; Bristol-Myers Squibb Co., P.O. Box 4000, Princeton, NJ 08543 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: HETEROCYCLIC CONTAINING BIPHENYL aP2 INHIBITORS AND METHOD</p> <div style="text-align: center; margin: 20px 0;">  <span style="margin: 0 20px;">(I)</span>  <span style="margin: 0 20px;">(a)</span> </div> <p>(57) Abstract</p> <p>aP2 inhibiting compounds are provided having formula (I) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, X-Z and (a) are as described herein. A method is also provided for treating diabetes and related diseases, especially Type II diabetes, employing such aP2 inhibitor or a combination of such aP2 inhibitor and another antidiabetic agent such as metformin, glyburide, troglitazone and/or insulin.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HETEROCYCLIC CONTAINING BIPHENYL aP2  
INHIBITORS AND METHOD

Field of the Invention

5           The present invention relates to heterocyclic  
containing biphenyls which are inhibitors of aP2 and to a  
method for treating diabetes, especially Type II  
diabetes, as well as hyperglycemia, hyperinsulinemia,  
10   obesity, Syndrome X, diabetic complications,  
atherosclerosis and related diseases, and other chronic  
inflammatory and autoimmune/inflammatory diseases,  
employing such heterocyclic containing biphenyls alone or  
in combination with one or more types of antidiabetic  
agents.

15

Background of the Invention

Fatty acid binding proteins (FABPs) are small  
cytoplasmic proteins which bind to fatty acids such as  
oleic acids which are important metabolic fuels and  
20   cellular regulators. Dysregulation of fatty acid  
metabolism in adipose tissue is a prominent feature of  
insulin resistance and the transition from obesity to  
non-insulin dependent diabetes mellitus (NIDDM or Type II  
diabetes).

25           aP2 (adipocyte fatty binding protein), an abundant  
14.6 KDa cytosolic protein in adipocytes, and one of a  
family of homologous intracellular fatty acid binding  
proteins (FABPs), is involved in the regulation of fatty  
acid trafficking in adipocytes and mediates fatty acid  
30   fluxes in adipose tissue. G.S. Hotamisligil et al,  
"Uncoupling of Obesity from Insulin Resistance Through a  
Targeted Mutation in aP2, the Adipocyte Fatty Acid  
Binding Protein", Science, Vol. 274, Nov. 22, 1996, pp.  
1377-1379, report that aP2-deficient mice placed on a  
35   high fat diet for several weeks developed dietary  
obesity, but, unlike control-mice on a similar diet, did  
not develop insulin resistance or diabetes. Hotamisligil


et al conclude that "aP2 is central to the pathway that links obesity to insulin resistance" (Abstract, page 1377).

DIALOG ALERT DBDR928 dated January 2, 1997,  
 5 Pharmaprojects No. 5149 (Knight-Ridder Information) discloses that a major drug company "is using virtual screening techniques to identify potential new antidiabetic compounds." It is reported that "the company is screening using aP2, a protein related to  
 10 adipocyte fatty acid binding protein."

U.S. application Serial No. 60/100,677, filed September 17, 1998 (attorney file LA24\*) discloses a method for treating diabetes employing an aP2 inhibitor.

15 Description of the Invention

In accordance with the present invention, heterocyclic containing biphenyl compounds are provided which have the structure



20 where  $R^1$  and  $R^2$  are the same or different and are independently selected from H, alkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heteroarylalkyl, aralkyl, cycloheteroalkyl and cycloheteroalkylalkyl;


$R^3$  is selected from hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkylalkyl, cycloalkenyl, alkylcarbonyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkenylalkyl, haloalkyl, polyhaloalkyl, cyano, nitro, hydroxy, amino, alkanoyl, alkylthio, alkylsulfonyl, alkoxycarbonyl,  
 25 alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyloxy, alkylaminosulfonyl, alkylamino, dialkylamino, all optionally substituted through available carbon atoms with 1, 2, 3, 4 or 5 groups selected from hydrogen, halo,  
 30


alkyl, polyhaloalkyl, alkoxy, haloalkoxy, polyhaloalkoxy,  
alkoxycarbonyl, alkenyl, alkynyl, cycloalkyl,  
cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl,  
hydroxy, hydroxyalkyl, nitro, cyano, amino, substituted  
5 amino, alkylamino, dialkylamino, thiol, alkylthio,  
alkylcarbonyl, acyl, alkoxycarbonyl, aminocarbonyl,  
alkynylaminocarbonyl, alkylaminocarbonyl,  
alkenylaminocarbonyl, alkylcarbonyloxy,  
alkylcarbonylamino, alkoxycarbonylamino, alkylsulfonyl,  
10 aminosulfinyl, aminosulfonyl, alkylsulfinyl, sulfonamido  
or sulfonyl;


R<sup>4</sup> is selected from hydrogen, halogen, alkyl,  
alkenyl, alkynyl, alkoxy, aryl, heteroaryl, arylalkyl,  
heteroarylalkyl, arylalkenyl, arylalkynyl, cycloalkyl,  
15 cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl,  
cycloalkenyl, cycloalkynyl, alkylcarbonyl, arylcarbonyl,  
cycloheteroalkyl, cycloheteroalkylalkyl,  
cycloalkenylalkyl, polycycloalkenyl,  
polycycloalkenylalkyl, polycycloalkynyl,  
20 polycycloalkynylalkyl, haloalkyl, polyhaloalkyl, cyano,  
nitro, hydroxy, amino, alkanoyl, aroyl, alkylthio,  
alkylsulfonyl, arylsulfonyl, alkoxycarbonyl,  
aryloxycarbonyl, alkylaminocarbonyl, arylaminocarbonyl,  
alkylcarbonylamino, alkylcarbonyloxy, alkylaminosulfonyl,  
25 arylaminosulfonyl, alkylamino, dialkylamino, all  
optionally substituted through available carbon atoms  
with 1, 2, 3, 4 or 5 groups selected from hydrogen, halo,  
alkyl, haloalkyl, polyhaloalkyl, alkoxy, haloalkoxy,  
polyhaloalkoxy, alkoxycarbonyl, alkenyl, alkynyl,  
30 cycloalkyl, cycloalkylalkyl, cycloheteroalkyl,  
cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl,  
arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy,  
aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxy,  
heteroarylalkyl, heteroarylalkenyl, heteroaryloxy,  
35 hydroxy, hydroxyalkyl, nitro, cyano, amino, substituted  
amino, alkylamino, dialkylamino, thiol, alkylthio,  
arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl,



- arylcarbonyl, acyl, arylaminocarbonyl, alkoxycarbonyl,  
 aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl,  
 alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy,  
 alkylcarbonylamino, arylcarbonylamino,  
 5 alkoxycarbonylamino, arylsulfinyl, arylsulfinylalkyl,  
 arylsulfonyl, alkylsulfonyl, aminosulfinyl,  
 aminosulfonyl, arylsulfonylamino,  
 heteroarylcarbonylamino, heteroarylsulfinyl,  
 heteroarylthio, heteroarylsulfonyl, alkylsulfinyl,  
 10 sulfonamido or sulfonyl;



X is a bond or a linker group selected from  $(CH_2)_n$ ,  
 $O(CH_2)_n$ ,  $S(CH_2)_n$ ,  $NHCO$ ,  $CH=CH$ , cycloalkylene or  
 $N(R^5)(CH_2)_n$ , (where  $n = 0-5$  and  $R^5$  is H, alkyl, or  
 alkanoyl);

- 15 Z is  $CO_2H$  or tetrazole of the formula  or  
 its tautomer; and



- the group  represents a heterocyclic group  
 (including heteroaryl and cycloheteroalkyl groups)  
 preferably containing 5-members within the ring and  
 20 containing preferably 1-3 heteroatoms within the ring,  
 and which may further optionally include one or two  
 substituents which are alkyl, alkenyl, hydroxyalkyl,  
 keto, carboxyalkyl, carboxy, cycloalkyl, alkoxy, formyl,  
 alkanoyl, alkoxyalkyl or alkoxycarboxyl;  
 25 with the provisos that (1)  $n \neq 0$  when Z is  $CO_2H$  and X  
 is  $O(CH_2)_n$ ,  $S(CH_2)_n$  or  $N(R^5)(CH_2)_n$ ; and



- (2) when  is , then X-Z may not be  
 O-lower alkylene- $CO_2H$  or -O-lower alkylene- $CO_2$ alkyl when  
 $R^1$  and  $R^2$  are both aryl or substituted aryl and  $R^3$  and  $R^4$   
 30 are each hydrogen;

and including pharmaceutically acceptable salts thereof, and prodrug esters thereof, and all stereoisomers thereof.

In addition, in accordance with the present invention, a method is provided for treating diabetes, especially Type II diabetes, and related diseases such as insulin resistance, hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis and other chronic inflammatory and autoimmune/inflammatory diseases, wherein a therapeutically effective amount of a compound of structure I (which inhibits  $\alpha P2$ ) is administered to a human patient in need of treatment.

The term "chronic inflammatory and autoimmune/inflammatory diseases" referred to above includes inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, rheumatoid arthritis, chronic obstructive pulmonary disease, emphysema, systemic lupus erythematosus, and other disease states involving tissue injury-, necrosis-, and/or infection-induced imbalanced inflammation associated with macrophage and leukocyte over-stimulation and excessive or dysregulated release of cellular mediators.

In addition, in accordance with the present invention, a method is provided for treating chronic and autoimmune/inflammatory diseases including inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, rheumatoid arthritis, chronic obstructive pulmonary disease, emphysema, systemic lupus erythematosus, and other disease states involving tissue injury-, necrosis-, and/or infection-induced imbalanced inflammation associated with macrophage and leukocyte over-stimulation and excessive or dysregulated release of cellular mediators, wherein a therapeutically effective amount of an  $\alpha P2$  inhibitor is administered to a human patient in need of treatment.

In addition to compounds of formula I, other aP2 inhibitors useful in carrying out the above method for treating chronic inflammatory and autoimmune/inflammatory diseases are disclosed in U.S. application Serial No. 09/390,275, filed September 7, 1999 (file LA24a), which is incorporated herein by reference.

The conditions, diseases, and maladies collectively referred to as "Syndrome X" (also known as Metabolic Syndrome) are detailed in Johannsson *J. Clin. Endocrinol. Metab.*, 82, 727-34 (1997).


The conditions, diseases and maladies collectively referred to as "diabetic complications" include retinopathy, neuropathy and nephropathy, and other known complications of diabetes.


In addition, in accordance with the present invention, a method is provided for treating diabetes and related diseases as defined above and hereinafter, as well as obesity, hypertriglyceridemia, Syndrome X, diabetic complications and other chronic inflammatory and autoimmune/inflammatory diseases, wherein a therapeutically effective amount of a combination of a compound of structure I and 1, 2, 3 or more other types of therapeutic agents is administered to a human patient in need of treatment.


The term "other type of therapeutic agents" as employed herein refers to one or more antidiabetic agents (other than aP2 inhibitors of formula I), one or more anti-obesity agents, one or more lipid-lowering agents (including anti-atherosclerosis agents), one or more anti-hypertensive agents, one or more anti-platelet agents, and/or one or more anti-infective agents.

In the above method of the invention, the compound of structure I will be employed in a weight ratio to the other type of therapeutic agent (depending upon its mode of operation) within the range from about 0.01:1 to about 500:1, preferably from about 0.1:1 to about 100:1.





Examples of the group  include (but are not limited to) heteroaryl groups and cycloheteroalkyl groups as defined herein and preferably include the following:




5

where  $R^8$  is selected from H, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, or alkenyl, and


$R^9$  and  $R^{9'}$  are the same or different and are selected independently from H, alkyl, alkoxy, alkenyl, formyl,  $CO_2H$ ,  $CO_2$ (lower alkyl), hydroxyalkyl, alkoxyalkyl,  $CO$ (alkyl), carboxylalkyl, haloalkyl, alkenyl or cycloalkyl.

With respect to the  $R^8$ ,  $R^9$  and  $R^{9'}$  groups, alkyl by itself or as part of another group will preferably contain 1 to 6 carbons.

Examples of X-Z moieties include (but are not limited to)



Preferred are compounds of formula I where





5

(where  $R^8$  is hydrogen, alkyl, fluoroalkyl or alkoxyalkyl, and where  $R^9$  is hydrogen, alkyl, fluoroalkyl, alkoxy or hydroxyalkyl).

10  $R^1$  and  $R^2$  are each phenyl, substituted phenyl or cycloalkyl;  $R^3$  and  $R^4$  are the same or different are independently selected from H, halo, alkyl or alkoxy; X is  $OCH_2$ ,  $NHCH_2$ ,  $CH_2$  or  $CH_2CH_2$ ; and Z is  $CO_2H$  or tetrazole.

More preferred are compounds of formula I where




(where R<sup>8</sup> is  
hydrogen, alkyl  
or fluoroalkyl)





(where R<sup>9</sup> is hydrogen,  
alkyl, fluoroalkyl  
or alkoxy)


- 5        R<sup>1</sup> and R<sup>2</sup> are each phenyl; R<sup>3</sup> and R<sup>4</sup> are each H; X  
is OCH<sub>2</sub>, CH<sub>2</sub> or NHCH<sub>2</sub>; and Z is CO<sub>2</sub>H or tetrazole.

#### Detailed Description of the Invention

- 10        Compounds of the invention of general structure I  
may be synthesized from intermediate II as shown in the  
schemes set out below. The groups R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup>, in  
intermediate II, are the same as described above with  
respect to the formula I compounds of the invention while  
A is a precursor to X-Z and is detailed below.









where  $\text{R}^{10}$  is lower alkyl or benzyl.


5 The biphenyl portion of the molecule may be prepared by reaction of compound III with substituted aryl IV via Stille or Suzuki type coupling to give compounds of the type V.







where W is  $\text{B}(\text{OH})_2$ ,  $\text{SnBu}_3$ , or  $\text{ZnBr}$  or  $\text{ZnCl}$  and G is Cl, Br, I, or OTf or G is  $\text{B}(\text{OH})_2$  or  $\text{SnBu}_3$  and W is Cl, Br, I, or OTf and E may be CHO, CN,  $\text{CO}_2\text{R}^{10}$ , OH,  $\text{N}(\text{R}^5)\text{H}$ ,  $\text{NO}_2$ ,  $\text{SR}^{10}$ ,  $\text{OR}^{10}$ ,  $\text{OSi}(\text{R}^{10})_3$ , or preferably X-Z or a protected variant thereof and where  $\text{R}^{11}$  is  $\text{CO}_2\text{R}^{10}$ , CHO, CN,  $-\text{NH}-\text{N}=\text{C}(\text{R}^2)(\text{CH}_2\text{R}^1)$ ,  $\text{NH}_2$ , or  $-\text{CONH}-\text{N}=\text{CH}(\text{R}^2)$


with the proviso that when W or G is ZnBr or ZnCl, E or R<sup>11</sup> may not be a reactive group such as CHO.



Compound V, depicted below where Y is , can be utilized to make heterocycles of the type VIA-VIN by standard methods described in the literature, for example, as shown below.








Alternately, and in some cases more preferably, compounds of the type VII may be converted to the desired  
 5 heterocycles by these aforementioned methodologies and subsequently converted to compounds of type II via biphenyl coupling reactions.






with the proviso that when W is  $\text{ZnCl}$  or  $\text{ZnBr}$ , E cannot be a reactive group such as  $\text{CHO}$ .

Unless otherwise indicated, the term "lower alkyl", "alkyl" or "alk" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons, containing 1 to 20 carbons, preferably 1 to 10 carbons, more preferably 1 to 8 carbons, in the normal chain, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethyl-pentyl, nonyl, decyl, undecyl, dodecyl, the various branched chain isomers thereof, and the like as well as such groups including 1 to 4 substituents such as halo, for example F, Br, Cl or I or  $\text{CF}_3$ , alkoxy, aryl, aryloxy, aryl(aryl) or diaryl, arylalkyl, arylalkyloxy, alkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkyloxy, amino, hydroxy, hydroxyalkyl, acyl, heteroaryl, heteroaryloxy, heteroarylalkyl, heteroarylalkoxy, aryloxyalkyl, aryloxyaryl, alkylamido, alkanoylamino, arylcarbonylamino, nitro, cyano, thiol, haloalkyl, trihaloalkyl and/or alkylthio and/or any of the  $\text{R}^3$  groups or substituents for  $\text{R}^3$ .

Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons

forming the rings, preferably 3 to 10 carbons, forming the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl,



any of which groups may be optionally substituted with 1 to 4 substituents such as halogen, alkyl, alkoxy, hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the R<sup>4</sup> groups or substituents for R<sup>4</sup>.

The term "cycloalkenyl" as employed herein alone or as part of another group refers to cyclic hydrocarbons containing 3 to 12 carbons, preferably 5 to 10 carbons and 1 or 2 double bonds. Exemplary cycloalkenyl groups include cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclohexadienyl, and cycloheptadienyl, which may be optionally substituted as defined for cycloalkyl.

The term "cycloalkylene" as employed herein refers to a "cycloalkyl" group which includes free bonds and thus is a linking group such as



and the like, and may optionally be substituted as defined above for "cycloalkyl".

The term "alkanoyl" as used herein alone or as part of another group refers to alkyl linked to a carbonyl group.

Unless otherwise indicated, the term "lower alkenyl" or "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons,

and more preferably 1 to 8 carbons in the normal chain, which include one to six double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, hydroxy, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonyl-amino, nitro, cyano, thiol, alkylthio and/or any of the R<sup>3</sup> groups, or the R<sup>3</sup> substituents set out herein.

Unless otherwise indicated, the term "lower alkynyl" or "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one triple bond in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonynyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, heteroaryl, cycloheteroalkyl, hydroxy, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, and/or alkylthio, and/or any of the R<sup>3</sup> groups, or the R<sup>3</sup> substituents set out herein.

The terms "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkenyl and alkynyl groups as described above having an aryl substituent.



Where alkyl groups as defined above have single bonds for attachment to other groups at two different


carbon atoms, they are termed "alkylene" groups and may optionally be substituted as defined above for "alkyl".

Where alkenyl groups as defined above and alkynyl groups as defined above, respectively, have single bonds  
 5 for attachment at two different carbon atoms, they are termed "alkenylene groups" and "alkynylene groups", respectively, and may optionally be substituted as defined above for "alkenyl" and "alkynyl".

Suitable alkylene, alkenylene or alkynylene groups  
 10  $(CH_2)_n$  or  $(CH_2)_p$  (where, p is 1 to 8, preferably 1 to 5, and n is 1 to 5, preferably 1 to 3, which includes alkylene, alkenylene or alkynylene groups) as defined herein, may optionally include 1, 2, or 3 substituents which include alkyl, alkenyl, halogen, cyano, hydroxy,  
 15 alkoxy, amino, thioalkyl, keto,  $C_3$ - $C_6$  cycloalkyl, alkylcarbonylamino or alkylcarbonyloxy.

Examples of  $(CH_2)_n$  or  $(CH_2)_p$ , alkylene, alkenylene and alkynylene include






The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine as well as CF<sub>3</sub>, with chlorine or fluorine being preferred.

The term "metal ion" refers to alkali metal ions such as sodium, potassium or lithium and alkaline earth metal ions such as magnesium and calcium, as well as zinc and aluminum.

Unless otherwise indicated, the term "aryl" as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings for example



and may be optionally substituted through available carbon atoms with 1, 2, or 3 groups selected from hydrogen, halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkyl-alkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, arylthio, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkyl-aminocarbonyl, arylaminocarbonyl, alkoxy carbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfon-aminocarbonyl and/or any of the R<sup>4</sup> groups or the R<sup>4</sup> substituents set out herein.

Unless otherwise indicated, the term "lower alkoxy", "alkoxy", "aryloxy" or "aralkoxy" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to an oxygen atom.

Unless otherwise indicated, the term "substituted amino" as employed herein alone or as part of another group refers to amino substituted with one or two substituents, which may be the same or different, such as

alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or thioalkyl. These substituents may be further substituted  
5 with a carboxylic acid and/or any of the R<sup>4</sup> groups or R<sup>4</sup> substituents thereof as set out above. In addition, the amino substituents may be taken together with the nitrogen atom to which they are attached to form  
1-pyrrolidinyl, 1-piperidinyl, 1-azepinyl, 4-morpholinyl,  
10 4-thiamorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, 1-pyrrolidinyl, 1-piperidinyl, or 1-azepinyl, optionally substituted with alkyl, alkoxy, alkylthio, halo, trifluoromethyl or hydroxy.

15 Unless otherwise indicated, the term "lower alkylthio", "alkylthio", "arylthio" or "aralkylthio" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to a sulfur atom.


20 Unless otherwise indicated, the term "lower alkylamino", "alkylamino", "arylamino", or "arylalkylamino" as employed herein alone or as part of another group includes any of the above alkyl, aryl or arylalkyl groups linked to a nitrogen atom.

25 Unless otherwise indicated, the term "acyl" as employed herein by itself or part of another group, as defined herein, refers to an organic radical linked to a

carbonyl  $\left( \begin{array}{c} \text{O} \\ \parallel \\ \text{C} \end{array} \right)$  group; examples of acyl groups include  
any of the R<sup>3</sup> groups attached to a carbonyl, such as  
30 alkanoyl, alkenoyl, aroyl, aralkanoyl, heteroaroyl, cycloalkanoyl, cycloheteroalkanoyl and the like.

Unless otherwise indicated, the term  
"cycloheteroalkyl" as used herein alone or as part of  
another group refers to a 5-, 6- or 7-membered saturated  
35 or partially unsaturated ring which includes 1 to 2 hetero atoms such as nitrogen, oxygen and/or sulfur, linked through a carbon atom or a heteroatom, where


possible, optionally via the linker  $(CH_2)_p$  (where  $p$  is 1, 2 or 3), such as




5

and the like. The above groups may include 1 to 4 substituents such as alkyl, halo, oxo and/or any of of the  $R^4$  groups, or the  $R^4$  substituents set out herein. In addition, any of the cycloheteroalkyl rings can be fused to a cycloalkyl, aryl, heteroaryl or cycloheteroalkyl ring.

Unless otherwise indicated, the term "heteroaryl" as used herein alone or as part of another group refers to a 5- or 6- membered aromatic ring which includes 1, 2, 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur, and such rings fused to an aryl, cycloalkyl, heteroaryl or cycloheteroalkyl ring (e.g. benzothiophenyl, indolyl), and includes possible N-oxides. The heteroaryl group may optionally include 1 to 4 substituents such as any of the  $R^4$  groups or the  $R^4$  substituents set out above. Examples of heteroaryl groups include the following:







5

and the like.

The term "cycloheteroalkylalkyl" as used herein alone or as part of another group refers to cycloheteroalkyl groups as defined above linked through a C atom or heteroatom to a  $(\text{CH}_2)_p$  chain.

10

The term "heteroarylalkyl" or "heteroarylalkenyl" as used herein alone or as part of another group refers to a heteroaryl group as defined above linked through a C atom or heteroatom to a  $-(\text{CH}_2)_p-$  chain, alkylene or alkenylene as defined above.

15

The term "polyhaloalkyl" as used herein refers to an "alkyl" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as  $\text{CF}_3\text{CH}_2$ ,  $\text{CF}_3$  or  $\text{CF}_3\text{CF}_2\text{CH}_2$ .

20


The term "polyhaloalkyloxy" as used herein refers to an "alkoxy" or "alkyloxy" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as  $\text{CF}_3\text{CH}_2\text{O}$ ,  $\text{CF}_3\text{O}$  or  $\text{CF}_3\text{CF}_2\text{CH}_2\text{O}$ .

25


The term "prodrug esters" as employed herein includes prodrug esters which are known in the art for carboxylic acids such as similar carboxylic acid esters such as methyl, ethyl benzyl and the like. Other examples include the following groups: (1-

30


alkanoyloxy)alkyl such as,




wherein  $\text{R}^a$ ,  $\text{R}^b$  and  $\text{R}^c$  are H, alkyl, aryl or aryl-alkyl; however  $\text{R}^a\text{O}$  cannot be HO, and where  $\text{Z}^1$  is





Examples of such prodrug esters include



10  $t\text{-C}_4\text{H}_9\text{CO}_2\text{CH}_2-$ , or



Other examples of suitable prodrug esters include



wherein  $\text{R}^a$  can be H, alkyl (such as methyl or t-butyl), arylalkyl (such as benzyl) or aryl (such as phenyl);  $\text{R}^d$  is H, alkyl, halogen or alkoxy,  $\text{R}^e$  is alkyl, aryl, 20 arylalkyl or alkoxy, and  $n_1$  is 0, 1 or 2.

Where the compounds of structure I are in acid form it may form a pharmaceutically acceptable salt such as alkali metal salts such as lithium, sodium or potassium, alkaline earth metal salts such as calcium or magnesium 25 as well as zinc or aluminum and other cations such as

ammonium, choline, diethanolamine, ethylenediamine, t-butylamine, t-octylamine, dehydroabietylamine.

5 All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one or the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures  
10 thereof. The processes for preparation can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional  
15 crystallization.

Where desired, the compounds of structure I may be used in combination with one or more other types of therapeutic agents which may be administered orally in the same dosage form, in a separate oral dosage form or  
20 by injection.

The other type of therapeutic agent which may be optionally employed in combination with the  $\alpha P2$  inhibitor of formula I may be 1,2,3 or more antidiabetic agents or antihyperglycemic agents including insulin secretagogues  
25 or insulin sensitizers, or other antidiabetic agents preferably having a mechanism of action different from  $\alpha P2$  inhibition and may include biguanides, sulfonyl ureas, glucosidase inhibitors, PPAR  $\gamma$  agonists, such as thiazolidinediones, SGLT2 inhibitors, PPAR  $\alpha/\gamma$  dual  
30 agonists, dipeptidyl peptidase IV (DP4) inhibitors, and/or meglitinides, as well as insulin, and/or glucagon-like peptide-1 (GLP-1).

It is believed that the use of the compounds of structure I in combination with 1, 2, 3 or more other  
35 antidiabetic agents produces antihyperglycemic results greater than that possible from each of these medicaments

alone and greater than the combined additive anti-hyperglycemic effects produced by these medicaments.

The other antidiabetic agent may be an oral antihyperglycemic agent preferably a biguanide such as metformin or phenformin or salts thereof, preferably metformin HCl.

Where the other antidiabetic agent is a biguanide, the compounds of structure I will be employed in a weight ratio to biguanide within the range from about 0.01:1 to about 100:1, preferably from about 0.1:1 to about 5:1.

The other antidiabetic agent may also preferably be a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the  $\beta$ -cells, with glyburide and glipizide being preferred, which may be administered in the same or in separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the sulfonyl urea in the range from about 0.01:1 to about 100:1, preferably from about 0.2:1 to about 10:1.

The oral antidiabetic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in the same or in a separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the glucosidase inhibitor within the range from about 0.01:1 to about 100:1, preferably from about 0.5:1 to about 50:1.

The compounds of structure I may be employed in combination with a PPAR  $\gamma$  agonist such as a thiazolidinedione oral anti-diabetic agent or other insulin sensitizers (which has an insulin sensitivity effect in NIDDM patients) such as troglitazone (Warner-

Lambert's Rezulin<sup>®</sup>, disclosed in U.S. Patent No. 4,572,912), rosiglitazone (SKB), pioglitazone (Takeda), Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016), Glaxo-Welcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer, isaglitazone (MIT/J&J), JTT-501 (JPNT/P&U), L-895645 (Merck), R-119702 (Sankyo/WL), NN-2344 (Dr. Reddy/NN), or YM-440 (Yamanouchi), preferably rosiglitazone and pioglitazone.

10           The compounds of structure I will be employed in a weight ratio to the thiazolidinedione in an amount within the range from about 0.01:1 to about 100:1, preferably from about 0.2:1 to about 10:1.

15           The sulfonyl urea and thiazolidinedione in amounts of less than about 150 mg oral antidiabetic agent may be incorporated in a single tablet with the compounds of structure I.

20           The compounds of structure I may also be employed in combination with a antihyperglycemic agent such as insulin or with glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492 to Habener, the disclosure of which is incorporated herein by reference), as well as AC2993 (Amylen) and LY-315902 (Lilly), which 25 may be administered via injection, intranasal, or by transdermal or buccal devices.

30           Where present, metformin, the sulfonyl ureas, such as glyburide, glimepiride, glipyrizide, glipizide, chlorpropamide and gliclazide and the glucosidase inhibitors acarbose or miglitol or insulin (injectable, pulmonary, buccal, or oral) may be employed in formulations as described above and in amounts and dosing as indicated in the Physician's Desk Reference (PDR).

35           Where present, metformin or salt thereof may be employed in amounts within the range from about 500 to about 2000 mg per day which may be administered in single or divided doses one to four times daily.

Where present, the thiazolidinedione anti-diabetic agent may be employed in amounts within the range from about 0.01 to about 2000 mg/day which may be administered in single or divided doses one to four times per day.

5       Where present insulin may be employed in formulations, amounts and dosing as indicated by the Physician's Desk Reference.

Where present GLP-1 peptides may be administered in oral buccal formulations, by nasal administration or  
10       parenterally as described in U.S. Patent Nos. 5,346,701 (TheraTech), 5,614,492 and 5,631,224 which are incorporated herein by reference.

The other antidiabetic agent may also be a PPAR  $\alpha/\gamma$  dual agonist such as AR-HO39242 (Astra/Zeneca), GW-  
15       409544 (Glaxo-Wellcome), KRP297 (Kyorin Merck) as well as those disclosed by Murakami et al, "A Novel Insulin Sensitizer Acts As a Coligand for Peroxisome Proliferation - Activated Receptor Alpha (PPAR alpha) and PPAR gamma. Effect on PPAR alpha Activation on Abnormal  
20       Lipid Metabolism in Liver of Zucker Fatty Rats", Diabetes 47, 1841-1847 (1998), and in U.S. provisional application No. 60/155,400, filed September 22, 1999, (attorney file LA29) the disclosure of which is incorporated herein by reference, employing dosages as set out therein, which  
25       compounds designated as preferred are preferred for use herein.

The other antidiabetic agent may be an SGLT2 inhibitor such as disclosed in U.S. provisional application No. 60/158,773, filed October 12, 1999  
30       (attorney file LA49), employing dosages as set out herein. Preferred are the compounds designated as preferred in the above application.

The other antidiabetic agent may be a DP4 inhibitor such as disclosed in WO99/38501, WO99/46272,  
35       WO99/67279 (PROBIODRUG), WO99/67278 (PROBIODRUG), WO99/61431 (PROBIODRUG), NVP-DPP728A (1-[[[2-[(5-cyanopyridin-2-yl)amino]ethyl]amino]acetyl]-2-cyano-(S)-

pyrrolidine) (Novartis) (preferred) as disclosed by Hughes et al, Biochemistry, 38(36), 11597-11603, 1999, TSL-225 (tryptophyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (disclosed by Yamada et al, Bioorg. & Med. Chem. Lett. 8 (1998) 1537-1540, 2-cyanopyrrolidides and 4-cyanopyrrolidides as disclosed by Ashworth et al, Bioorg. & Med. Chem. Lett., Vol. 6, No. 22, pp 1163-1166 and 2745-2748 (1996) employing dosages as set out in the above references.

10           The meglitinide which may optionally be employed in combination with the compound of formula I of the invention may be repaglinide, nateglinide (Novartis) or KAD1229 (PF/Kissei), with repaglinide being preferred.

15           The  $\alpha$ P2 inhibitor of formula I will be employed in a weight ratio to the meglitinide, PPAR  $\gamma$  agonist, PPAR  $\alpha/\gamma$  dual agonist, SGLT2 inhibitor or DP4 inhibitor within the range from about 0.01:1 to about 100:1, preferably from about 0.2:1 to about 10:1.


20           The hypolipidemic agent or lipid-lowering agent which may be optionally employed in combination with the compounds of formula I of the invention may include 1,2,3 or more MTP inhibitors, HMG CoA reductase inhibitors, squalene synthetase inhibitors, fibric acid derivatives, ACAT inhibitors, lipoxygenase inhibitors, cholesterol  
25           absorption inhibitors, ileal  $\text{Na}^+$ /bile acid cotransporter inhibitors, upregulators of LDL receptor activity, bile acid sequestrants, and/or nicotinic acid and derivatives thereof.

30           MTP inhibitors employed herein include MTP inhibitors disclosed in U.S. Patent No. 5,595,872, U.S. Patent No. 5,739,135, U.S. Patent No. 5,712,279, U.S. Patent No. 5,760,246, U.S. Patent No. 5,827,875, U.S. Patent No. 5,885,983 and U.S. Application Serial No. 09/175,180 filed October 20, 1998, now U.S. Patent No.  
35           5,962,440. Preferred are each of the preferred MTP inhibitors disclosed in each of the above patents and applications.

All of the above U.S. Patents and applications are incorporated herein by reference.

Most preferred MTP inhibitors to be employed in accordance with the present invention include preferred MTP inhibitors as set out in U.S. Patent Nos. 5,739,135 and 5,712,279, and U.S. Patent No. 5,760,246.

The most preferred MTP inhibitor is 9-[4-[4-[[2-(2,2,2-Trifluoroethoxy)benzoyl]amino]-1-piperidinyl]butyl]-N-(2,2,2-trifluoroethyl)-9H-fluorene-9-carboxamide



10

The hypolipidemic agent may be an HMG CoA reductase inhibitor which includes, but is not limited to, mevastatin and related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed in U.S. Patent Nos. 4,448,784 and 4,450,171. Other HMG CoA reductase inhibitors which may be employed herein include, but are not limited to, fluvastatin, disclosed in U.S. Patent No. 5,354,772, cerivastatin disclosed in U.S. Patent Nos. 5,006,530 and 5,177,080, atorvastatin disclosed in U.S. Patent Nos. 4,681,893, 5,273,995, 5,385,929 and 5,686,104, atavastatin (Nissan/Sankyo's nisvastatin (NK-104)) disclosed in U.S. Patent No. 5,011,930, Shionogi-Astra/Zeneca visastatin (ZD-4522) disclosed in U.S. Patent No. 5,260,440, and related statin compounds disclosed in U.S. Patent No. 5,753,675, pyrazole analogs



of mevalonolactone derivatives as disclosed in U.S. Patent No. 4,613,610, indene analogs of mevalonolactone derivatives as disclosed in PCT application WO 86/03488, 6-[2-(substituted-pyrrol-1-yl)-alkyl]pyran-2-ones and  
5 derivatives thereof as disclosed in U.S. Patent No. 4,647,576, Searle's SC-45355 (a 3-substituted pentanedioic acid derivative) dichloroacetate, imidazole analogs of mevalonolactone as disclosed in PCT application WO 86/07054, 3-carboxy-2-hydroxy-propane-  
10 phosphonic acid derivatives as disclosed in French Patent No. 2,596,393, 2,3-disubstituted pyrrole, furan and thiophene derivatives as disclosed in European Patent Application No. 0221025, naphthyl analogs of mevalonolactone as disclosed in U.S. Patent No.  
15 4,686,237, octahydronaphthalenes such as disclosed in U.S. Patent No. 4,499,289, keto analogs of mevinolin (lovastatin) as disclosed in European Patent Application No. 0,142,146 A2, and quinoline and pyridine derivatives disclosed in U.S. Patent No. 5,506,219 and 5,691,322.  
20 In addition, phosphinic acid compounds useful in inhibiting HMG CoA reductase suitable for use herein are disclosed in GB 2205837.

The squalene synthetase inhibitors suitable for use herein include, but are not limited to,  $\alpha$ -phosphono-  
25 sulfonates disclosed in U.S. Patent No. 5,712,396, those disclosed by Biller et al, J. Med. Chem., 1988, Vol. 31, No. 10, pp 1869-1871, including isoprenoid (phosphinyl-methyl)phosphonates as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S.  
30 Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., Current Pharmaceutical Design, 2, 1-40 (1996).

In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid  
35 pyrophosphates disclosed by P. Ortiz de Montellano et al, J. Med. Chem., 1977, 20, 243-249, the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-

PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc., 1976, 98, 1291-1293, phosphinylphosphonates reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544 and cyclopropanes reported by Capson, T.L., PhD  
5 dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

Other hypolipidemic agents suitable for use herein include, but are not limited to, fibric acid derivatives,  
10 such as fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine,  
15 colestipol and DEAE-Sephadex (Secholex®, Policexide®), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe  
20 Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid, acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin,  
25 poly(diallylmethylamine) derivatives such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

30 The other hypolipidemic agent may be an ACAT inhibitor such as disclosed in, Drugs of the Future 24, 9-15 (1999), (Avasimibe); "The ACAT inhibitor, Cl-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al,  
35 Atherosclerosis (Shannon, Irel). (1998), 137(1), 77-85; "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by

selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo, Cardiovasc. Drug Rev. (1998), 16(1), 16-30; "RP 73163: a bioavailable alkylsulfinyl-diphenylimidazole ACAT inhibitor", Smith, C., et al, Bioorg. Med. Chem. Lett. (1996), 6(1), 47-50; "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic activities in experimental animals", Krause et al, Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., Inflammation: Mediators Pathways (1995), 173-98, Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al, Curr. Med. Chem. (1994), 1(3), 204-25; "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. 6. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). 7. Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity", Stout et al, Chemtracts: Org. Chem. (1995), 8(6), 359-62, or TS-962 (Taisho Pharmaceutical Co. Ltd).

The hypolipidemic agent may be an upregulator of LD2 receptor activity such as MD-700 (Taisho Pharmaceutical Co. Ltd) and LY295427 (Eli Lilly).

The hypolipidemic agent may be a cholesterol absorption inhibitor preferably Schering-Plough's SCH48461 as well as those disclosed in Atherosclerosis 115, 45-63 (1995) and J. Med. Chem. 41, 973 (1998).

The hypolipidemic agent may be an ileal Na<sup>+</sup>/bile acid cotransporter inhibitor such as disclosed in Drugs of the Future, 24, 425-430 (1999).

Preferred hypolipidemic agents are pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin, atavastatin and ZD-4522.

The above-mentioned U.S. patents are incorporated herein by reference. The amounts and dosages employed

will be as indicated in the Physician's Desk Reference and/or in the patents set out above.

The compounds of formula I of the invention will be employed in a weight ratio to the hypolipidemic agent  
5 (were present), within the range from about 500:1 to about 1:500, preferably from about 100:1 to about 1:100.

The dose administered must be carefully adjusted according to age, weight and condition of the patient, as well as the route of administration, dosage form and  
10 regimen and the desired result.

The dosages and formulations for the hypolipidemic agent will be as disclosed in the various patents and applications discussed above.

The dosages and formulations for the other  
15 hypolipidemic agent to be employed, where applicable, will be as set out in the latest edition of the Physicians' Desk Reference.

For oral administration, a satisfactory result may be obtained employing the MTP inhibitor in an  
20 amount within the range of from about 0.01 mg/kg to about 500 mg and preferably from about 0.1 mg to about 100 mg, one to four times daily.

A preferred oral dosage form, such as tablets or capsules, will contain the MTP inhibitor in an amount of  
25 from about 1 to about 500 mg, preferably from about 2 to about 400 mg, and more preferably from about 5 to about 250 mg, one to four times daily.

For oral administration, a satisfactory result may be obtained employing an HMG CoA reductase inhibitor, for  
30 example, pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin in dosages employed as indicated in the Physician's Desk Reference, such as in an amount within the range of from about 1 to 2000 mg, and preferably from about 4 to about 200 mg.

35 The squalene synthetase inhibitor may be employed in dosages in an amount within the range of from about 10

mg to about 2000 mg and preferably from about 25 mg to about 200 mg.

5 A preferred oral dosage form, such as tablets or capsules, will contain the HMG CoA reductase inhibitor in an amount from about 0.1 to about 100 mg, preferably from about 5 to about 80 mg, and more preferably from about 10 to about 40 mg.

10 A preferred oral dosage form, such as tablets or capsules will contain the squalene synthetase inhibitor in an amount of from about 10 to about 500 mg, preferably from about 25 to about 200 mg.

The other hypolipidemic agent may also be a lipoxxygenase inhibitor including a 15-lipoxxygenase (15-LO) inhibitor such as benzimidazole derivatives as disclosed in WO 97/12615, 15-LO inhibitors as disclosed in WO 97/12613, isothiazolones as disclosed in WO 96/38144, and 15-LO inhibitors as disclosed by Sendobry et al "Attenuation of diet-induced atherosclerosis in rabbits with a highly selective 15-lipoxxygenase inhibitor lacking significant antioxidant properties, Brit. J. Pharmacology (1997) 120, 1199-1206, and Cornicelli et al, "15-Lipoxxygenase and its Inhibition: A Novel Therapeutic Target for Vascular Disease", Current Pharmaceutical Design, 1999, 5, 11-20.

25 The compounds of formula I and the hypolipidemic agent may be employed together in the same oral dosage form or in separate oral dosage forms taken at the same time.

30 The compositions described above may be administered in the dosage forms as described above in single or divided doses of one to four times daily. It may be advisable to start a patient on a low dose combination and work up gradually to a high dose combination.

35 The preferred hypolipidemic agent is pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin or cerivastatin.

The other type of therapeutic agent which may be optionally employed with the aP2 inhibitor of formula I may be 1, 2, 3 or more of an anti-obesity agent including a beta 3 adrenergic agonist, a lipase inhibitor, a  
5 serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta drug and/or an anorectic agent.

The beta 3 adrenergic agonist which may be optionally employed in combination with a compound of formula I may be AJ9677 (Takeda/Dainippon), L750355  
10 (Merck), or CP331648 (Pfizer) or other known beta 3 agonists as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064, with AJ9677, L750,355 and CP331648 being preferred.

The lipase inhibitor which may be optionally  
15 employed in combination with a compound of formula I may be orlistat or ATL-962 (Alizyme), with orlistat being preferred.

The serotonin (and dopamine) reuptake inhibitor which may be optionally employed in combination with a  
20 compound of formula I may be sibutramine, topiramate (Johnson & Johnson) or axokine (Regeneron), with sibutramine and topiramate being preferred.

The thyroid receptor beta compound which may be optionally employed in combination with a compound of  
25 formula I may be a thyroid receptor ligand as disclosed in WO97/21993 (U. Cal SF), WO99/00353 (KaroBio) and GB98/284425 (KaroBio), with compounds of the KaroBio applications being preferred.

The anorectic agent which may be optionally  
30 employed in combination with a compound of formula I may be dexamphetamine, phentermine, phenylpropanolamine or mazindol, with dexamphetamine being preferred.

The various anti-obesity agents described above may be employed in the same dosage form with the compound of  
35 formula I or in different dosage forms, in dosages and regimens as generally known in the art or in the PDR.

The other type of therapeutic agent which may be optionally employed with the  $\text{aP2}$  inhibitor of formula I may be 1, 2, 3 or more of an antihypertensive agent including an ACE inhibitor, a vasopeptidase inhibitor, an angiotensin II antagonist, a calcium channel blocker, a potassium channel opener, an  $\alpha$ -blocker, a  $\beta$  blocker, a centrally acting  $\alpha$  agonist, and/or a diuretic.

The ACE inhibitor which may be optionally employed in combination with a compound of formula I may be lisinopril, enalapril, quinapril, benazepril, fosinopril, fentiapril, ramipril, captopril, enalaprilat, moexipril, tranolapril, perindopril, ceranopril, zofenopril or cetapril.

Preferred ACE inhibitors are captopril, as well as fosinopril, enalapril, lisinopril, quinapril, benazepril, fentiapril, ramipril, and moexipril.

The vasopeptidase inhibitor (also known as NEP/ACE inhibitors) which may be optionally employed with the  $\text{aP2}$  inhibitor of formula I may be omapatrilat (most preferred) and [S-(R\*,R\*)]-hexahydro-6-[(2-mercapto-1-oxo-3-phenylpropyl)amino]-2,2-dimethyl-7-oxo-1H-azepine-1-acetic acid (BMS 189,921 also preferred), as well as those disclosed in U.S. Patent Nos. 5,362,727, 5,366,973, 5,225,401, 4,722,810, 5,223,516, 4,749,688. U.S. Patent No. 5,504,080, U.S. Patent No. 5,552,397, U.S. Patent No. 5,612,359, U.S. Patent No. 5,525,723, European Patent Application 0599,444, 0481,522, 0599,444, 0595,610, European Patent Application 0534363.A2, 534,396 and 534,492, and European Patent Application 0629627.A2.

Preferred are those NEP/ACE inhibitors which are designated as preferred in the above patents/applications which U.S. patents/applications are incorporated herein by reference.

The angiotensin II receptor antagonist (also referred to herein as angiotensin II antagonist or AII antagonist) which may be optionally employed in

combination with a compound of formula I may be irbesartan, losartan, valsartan, candesartan, telmisartan, tasosartan and/or eprosartan, with irbesartan or losartan being preferred.

5           The calcium channel blocker (also referred to as a calcium antagonist) which may be optionally employed in combination with a compound of formula I may be amlodipine, diltiazem, nifedipine, verapamil, feldodipine, nisoldipine, isradipine and/or nicardipine,  
10 with amlodipine, diltiazem, verapamil and nifedipine being preferred.

          The alpha-blocker which may be optionally employed in combination with a compound of formula I may be terazosin, doxazosin or prazosin, all of which are  
15 preferred.

          The beta-blocker which may be optionally employed in combination with a compound of formula I may be nadolol, atenolol, propranolol, metoprolol, carvediol or sotalol, with atenolol and nadolol being preferred.

20           The potassium channel opener which may be optionally employed in combination with a compound of formula I may be minoxidil.

          The centrally acting  $\alpha$  agonist antihypertensive agent which may be optionally employed in combination  
25 with a compound of formula I may be clonidine or guanfacine, with clonidine being preferred.

          The diuretic which may be optionally employed in connection with a compound of formula I may be hydrochlorothiazide, torasemide, furosemide,  
30 spironolactone and/or indapamide, with hydrochlorothiazide and furosemide being preferred.

          The antiplatelet agent (also known as platelet aggregation inhibitor) which may be optionally employed in combination with a compound of formula I may be  
35 aspirin, clopidogrel, ticlopidine, dipyridamole, abciximab, tirofiban, eptifibatide, anagrelide and/or ifetroban, with aspirin and clopidogrel being preferred.



The anti-infective agent which may be optionally employed in combination with a compound of formula I may be an anti-infective that is effective against chlamydial infections, such as azithromycin, gatifloxacin, ciprofloxacin, levofloxacin and trovafloxacin, with azithromycin and gatifloxacin being preferred.

The various antihypertensive agents and antiplatelet agents and anti-infective agents described above may be employed in the same dosage form with the compound of formula I or in different dosage forms, in dosages and regimens as generally known in the art or in the PDR.

In carrying out the method of the invention, a pharmaceutical composition will be employed containing the compounds of structure I, with or without another therapeutic agent, in association with a pharmaceutical vehicle or diluent. The pharmaceutical composition can be formulated employing conventional solid or liquid vehicles or diluents and pharmaceutical additives of a type appropriate to the mode of desired administration. The compounds can be administered to mammalian species including humans, monkeys, dogs, etc. by an oral route, for example, in the form of tablets, capsules, granules or powders, or they can be administered by a parenteral route in the form of injectable preparations. The dose for adults is preferably between 20 and 2,000 mg per day, which can be administered in a single dose or in the form of individual doses from 1-4 times per day.

A typical capsule for oral administration contains compounds of structure I (250 mg), lactose (75 mg) and magnesium stearate (15 mg). The mixture is passed through a 60 mesh sieve and packed into a No. 1 gelatin capsule.

A typical injectable preparation is produced by aseptically placing 250 mg of compounds of structure I into a vial, aseptically freeze-drying and sealing. For use, the contents of the vial are mixed with 2 mL of

physiological saline, to produce an injectable preparation.

aP2 inhibitor activity of the compounds of the invention may be determined by use of an in vitro assay system which measures the potentiation of inhibition of aP2 by displacement of a fluorescent substrate from aP2 by the inhibitor. Inhibition constants ( $K_i$  values) for the aP2 inhibitors of the invention may be determined by the method described below:

10

**Production of purified recombinant human aP2 protein.**

Recombinant human aP2 protein is produced by standard recombinant DNA technology. In the typical case, aP2 is produced by heterologous expression in E. coli strain BL21(D53) transformed with pET11a vector containing the full length human aP2 cDNA (Baxa, C.A., Sha, R.S., Buelte, M.K., Smith, A.J., Matarese, V., Chinander, L.L., Boundy, K.L., and Bernlohr, D.A. (1989). Human adipocyte lipid-binding protein: purification of the protein and cloning of its complementary DNA. Biochemistry 28: 8683-8690 and Xu, Z., Buelte, M.K., Banaszak, L.J., and Bernlohr, D.A. (1991). Expression, purification and crystallization of the adipocyte lipid binding protein. J. Biol. Chem. 266: 14367-14370). Purification of aP2 from E. coli is conducted as described by Xu, yielding essentially homogeneous aP2 protein with molecular weight ~14600 daltons and free of endogenous fatty acids. The purified aP2 is capable of binding up to one mole of free fatty acid per mole protein. The binding and structural properties of recombinant aP2 protein were previously shown to be identical to aP2 protein isolated from adipose tissue.

**In vitro assay of aP2 inhibitors.** Inhibitors of aP2 are evaluated in a homogeneous fluorescent-based competition assay using recombinant aP2 protein and 1,8-anilino-naphthalene-sulfonic acid (1,8-ANS) as assay substrate.

This competition assay was adapted from generalized procedures described previously (Kane, C.D. and Bernlohr, D.A. (1996). A simple assay for intracellular lipid-binding proteins using displacement of 1-anilino-8-sulfonic acid. (1996) Anal. Biochem. 233: 197-204 and Kurian E., Kirk, W.R. and Prendergast, F.G. (1996) Affinity of fatty acid for r-rat intestinal fatty acid binding protein. Biochemistry, 35, 3865-3874). The method relies on the increase in fluorescence quantum yield of 1,8-ANS upon binding to the fatty acid binding site of aP2. The assay is run using appropriate concentrations of inhibitor, 1,8-ANS, and aP2 protein, in order to calculate the inhibitor binding constant ( $K_i$ ) for compounds being evaluated. The  $K_i$  calculation was based on the procedure previously described for calculation of dissociation constants described by Kurian. Lower  $K_i$  values indicate higher affinities of compounds binding to aP2.


In the assay as conducted for the inhibitors described herein, a series of aliquots of aP2 (5  $\mu$ M) in solution in 10 mM potassium phosphate buffer (pH 7.0) are mixed with an equimolar concentration of test compound, followed by the addition of a series of increasing concentrations of 1,8-ANS (from 0 to 5  $\mu$ M). The assay typically is conducted in 96-well plate format with reagents added using robotic instrumentation (Packard Multiprobe 104). The fluorescence value for each test is determined using a Cytofluor-4000 multi-well fluorescence plate reader (Perceptive Biosystems) using excitation wavelength 360 nm and emission wavelength 460 nm, or using other suitable spectrofluorometer. In preparation for the assay, test compounds are initially prepared at 10 mM in dimethylsulfoxide. All subsequent dilutions and assay additions are made in 10 mM potassium phosphate buffer, pH 7.0.

X-ray crystallography of the inhibitor-aP2 complex can be performed by one skilled in the art using


contemporary biophysical methodologies and commercial instrumentation. Such crystallographic data can be used to conclusively determine if a compound used in the present invention has embodied the structural requirement  
5 necessary for inhibition of aP2. An example of such an X-ray crystallographic determination is presented below:

Crystals of aP2 complexed with the inhibitors were typically grown by the hanging drop method. aP2, at 8.3 mg/ml, was pre-equilibrated with 1-5 mM of the inhibitor  
10 in 0.1 M Tris-HCl pH 8.0, 1% w/v DMSO for four hours. 2  $\mu$ l drops containing equilibrated protein and reservoir solution at a 1:1 ratio were suspended on plastic cover slips and equilibrated against a 1 ml reservoir containing 2.6-3.0 M ammonium sulfate in 0.1 M Tris-HCl  
15 pH 8.0. Crystals typically appeared in 2-3 days and reached maximum size within 2 weeks. Data was typically collected on a single flash-frozen crystal (Oxford Cryosystems) using a Rigaku rotating anode and an R-axis II image plate detector of a Bruker multiwire area  
20 detector. Diffraction from aP2 crystals was excellent. Diffraction was consistently observed to better than 2.0 Å resolution often to beyond 1.5 Å resolution. Data was processed either with DENZO/SCALEPACK (R-axis II data), or Xengen (Bruker data). XPLOR was used for structure  
25 refinement and model building was done using the molecular modeling package CHAIN. After a single round of refinement, examination of the  $F_o - F_c$  map typically allowed facile building of the inhibitor into aP2 binding cavity. Iterative fitting and refinement were continued  
30 until improvement was no longer seen in the electron density map or R-free.

The following working Examples represent preferred embodiments of the invention.

Example 1

A.




5

A solution of 2-bromo-benzoic acid (21.8 g, 108 mmol), 4-dimethyl amino pyridine (1 g, 8.19 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (22.6 g, 118 mmol) and benzoin (20 g, 94.2 mmol) in dichloromethane (200 mL) was stirred at room temperature for 4 hr. The reaction mixture was washed with water (2x50 mL), brine (50 mL), dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the ketoester as a colorless oil (36 g).


15 A mixture of ketoester (36 g, 94.2 mmol) and ammonium acetate (36 g, 471 mmol) in glacial acetic acid (350 mL) was stirred at reflux for 1.5 hr. The reaction mixture was cooled, diluted with water, and extracted with ethyl acetate (3x300 mL). The combined organic  
20 layers were washed with water (2x100 mL), saturated aqueous sodium bicarbonate solution (2x100 mL), and brine (100 mL), dried over anhydrous  $\text{MgSO}_4$  and concentrated to get yellow-orange oil. The crude product was  
25 recrystallized from hot methanol to get the title compound as off-white solid (21.6 g, 61%).

B.



Nitrogen was bubbled through a solution of Part A  
5 compound (1.48 g, 3.93 mmol), 3-formyl phenyl boronic  
acid (766 mg, 5.11 mmol) and aqueous sodium carbonate  
(3.2 mL, 2M, 6.4 mmol) in toluene (10 mL) and ethanol  
(4.2 mL) at room temperature for 30 min.  
Tetrakis(triphenylphosphine) palladium(0) (150 mg, 0.13  
10 mmol) was added and the mixture was stirred at 80°C for  
14 hr under argon. The reaction mixture was cooled to  
room temperature and extracted with ethyl acetate. The  
combined organic extracts were washed with brine, dried  
over anhydrous MgSO<sub>4</sub> and concentrated. The crude product  
15 was purified by flash chromatography on silica gel,  
eluting with a step gradient of 1:9 to 2:8 ethyl  
acetate/hexane to give title compound as a colorless oil  
(800 mg, 51%).


20 C.



Sodium borohydride (19 mg, 0.5 mmol) was added  
slowly to a solution of Part B compound (494 mg, 1.23  
25 mmol) in anhydrous methanol (2 mL) at 0°C. The reaction  
was stirred for 2 hr at 0°C, and then warmed to room  
temperature. The pH was adjusted to 2 with conc. HCl and  
concentrated. The residue was partitioned between ethyl  
acetate and water. The organic layer was washed with

water, dried ( $\text{MgSO}_4$ ) and concentrated to give the title compound as a white foam (360 mg, 89%).

D.




5

To a stirred solution of Part C compound (7.5 g, 18.6 mmol) in dioxane (62 mL) at room temperature was added zinc chloride (78 mg, 0.54 mmol) followed by thionyl chloride (2.74 mL, 37.5 mmol). The reaction was stirred for 1 hr and concentrated. The residue was dissolved in ethyl acetate, washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the title compound as a viscous oil (8.0 g, 100%).


15

E.




A solution of Part D compound (8.0 g, 18.6 mmol) and sodium cyanide (1.39 g, 28.2 mmol) in acetonitrile (75 mL) and water (7 mL) was stirred at reflux for 14 hr. The reaction mixture was cooled to room temperature, concentrated. The residue was partitioned between dichloromethane and water. The organic layer was dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 5:95 to 15:85 ethyl acetate/hexane to give title compound as a white solid (6.4 g, 83.6%).

F.




A solution of Part E compound (6.18 g, 15.0 mmol)  
5 and azidotrimethyltin (4.32 g, 21.0 mmol) in xylene (50 mL) was refluxed for 18 hr. The reaction mixture was cooled to room temperature, methanol (150 mL) was added and the mixture was stirred for 30 min and concentrated. The crude product was purified by flash chromatography on  
10 silica gel, eluting with a step gradient of 3:7 to 5:5 ethyl acetate/hexane and then with 95:5 dichloromethane/methanol to give title compound as a white solid (5.96 g, 87%). The compound can be recrystallized from hot ethanol.

15

Example 2

A.



20 Example 1 Part B compound (401 mg, 1 mmol) was added to a solution of silicon tetrachloride (170 mg, 115  $\mu$ L, 1 mmol) and sodium azide (195 mg, 3 mmol) in anhydrous




acetonitrile (3 mL) at 0°C. The reaction mixture was warmed to room temperature, then to 50°C and stirred for 24 hr. The reaction was cooled to room temperature and poured into aqueous saturated sodium carbonate solution.

5 The pH of the mixture was adjusted between 9-10 with sodium carbonate solution and it was extracted with chloroform (3x10 mL). The combined organic layers were washed with water, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash

10 chromatography on silica gel, eluting with a step gradient of 3:97 to 5:95 ethyl acetate/hexane to give title compound as a colorless oil (200 mg, 50%).

B.



15

A solution of Part A compound (199 mg, 0.5 mmol) and azidotrimethyltin (145 mg, 0.7 mmol) in xylene (5 mL) was heated at 100°C for 18 hr. The reaction mixture was


20 cooled to room temperature, methanol (8 mL) was added, stirred for 30 min, filtered and concentrated. Flash chromatography on silica gel, eluting with 1:4 ethyl acetate/hexane and then with 95:5:0.15

dichloromethane/methanol/acetic acid gave impure product.


25 Further purification by flash chromatography on silica gel, eluting with a step gradient of 1:4 to 1:1 ethyl acetate/hexane and then with 95:5 to 90:10

dichloromethane/methanol gave the title compound as a foam (70 mg, 32%).

30


Example 3

A.




- 5 Nitrogen was bubbled through a solution of Example 1 Part A compound (5 g, 13.29 mmol), 3-amino phenyl boronic acid (2.37 g, 17.29 mmol) and sodium carbonate (11 mL, 2M, 22 mmol) in toluene (40 mL) and ethanol (15 mL) at room temperature for 30 min.
- 10 Tetrakis(triphenylphosphine)-palladium(0) (508 mg, 0.43 mmol) was added and the reaction mixture was heated at 80°C for 18 hr. The reaction mixture was cooled to room temperature and partitioned between diethyl ether and water. The organic layer was washed with water, brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography eluting with 1:3 ethyl acetate/hexane to give title compound as a pale yellow foam (2.56 g, 49%).

20 B.




To a slurry of Part A compound (2.56 g, 6.60 mmol) and potassium carbonate (918 mg, 6.60 mmol) in anhydrous dimethyl formamide (20 mL) was added *tert*-butyl bromoacetate (975  $\mu$ L, 6.60 mmol). The reaction mixture  
5 was stirred at room temperature for 18 hr. Additional *tert*-butyl bromoacetate (150  $\mu$ L, 1.01 mmol) was added and the mixture was stirred for 7 hr. The reaction was partitioned between diethyl ether and aqueous sodium bicarbonate solution. The organic layer was washed with  
10 water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 15:85 ethyl acetate/hexane to give title compound as a foam (2.230 g, 67%).


15 C.




A solution of Part B compound (2.230 g, 4.4 mmol) in trifluoroacetic acid (15 mL) and dichloromethane (40 mL) was stirred at room temperature for 18 hr. The reaction  
20 mixture was concentrated, the residue was partitioned between ethyl acetate and water. The aqueous layer was basified with 1N NaOH and then the pH was adjusted between 3-4 with 10% citric acid solution. The organic layer was separated and washed with water and brine,  
25 dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 7:93 methanol/dichloromethane to give an oily foam which was triturated from ethyl acetate/hexane to get the title compound as a yellow foam (1.692 g, 86%).

Example 4

To a solution of Example 3 Part A compound (149 mg, 0.38 mmol) in dichloromethane (2 mL) was added diisopropylethylamine (100  $\mu$ L), followed by ethyl oxalyl chloride (147  $\mu$ L, 0.42 mmol). The reaction was stirred for 2 hr, partitioned between ethyl acetate and aqueous HCl (1N). The organic layer was washed with water, saturated sodium bicarbonate solution, and brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to get the crude ester as a yellow-brown foam (189 mg). To a solution of the crude ester in methanol (3 mL) was added dropwise aqueous NaOH (~800  $\mu$ L, 1N) until the solution became turbid. Precipitate was formed after stirring the solution overnight. The reaction was partitioned between ethyl acetate and aqueous HCl (1N). The organic layer was washed with water, and brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was triturated with ethyl acetate and hexane to give the title compound as an off-white solid (118 mg, 67%).


Example 5

A.




A mixture of benzil (10 g, 47.6 mmol) and ammonium  
5 acetate (44 g, 564 mmol) in glacial acetic acid (260 mL)  
was treated with 2-bromobenzaldehyde (5.5 mL, 47.6 mmol)  
and stirred at 105°C for 3 hr. The reaction mixture was  
cooled, diluted with water, and the precipitate was  
filtered. The solid was dissolved in warm ethyl acetate.  
10 The organic layer was washed with water, and brine, dried  
over anhydrous MgSO<sub>4</sub> and concentrated. The crude product  
was triturated from ethyl acetate/hexane to give the  
title compound as an off-white solid (16.55 g, 93%).

15 B.




To a solution of Part A compound (2.5 g, 6.67 mmol)  
and ethyl iodide (1.15 g, 7.3 mmol) in anhydrous  
20 dimethylformamide (17 mL) was added potassium carbonate  
(922 mg, 6.67 mmol). The reaction mixture was stirred at  
room temperature overnight, partitioned between ethyl  
acetate/water. The organic layer was washed with water,  
dried over anhydrous MgSO<sub>4</sub> and concentrated to give the  
25 title compound as a semi-solid (2.5 g, 94%).

C.



- 5 Nitrogen was bubbled through a solution of Part B compound (2.0 g, 4.96 mmol), 3-formyl phenyl boronic acid (966 mg, 6.44 mmol) and aqueous sodium carbonate (4 mL, 2M, 8 mmol) in toluene (15 mL) and ethanol (8 mL) at room temperature for 30 min. Tetrakis(triphenylphosphine)
- 10 palladium(0) (190 mg, 0.16 mmol) was added and the mixture was stirred at 80°C for 36 hr under argon. The reaction mixture was cooled to room temperature and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous
- 15 MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 1:9 to 2:8 ethyl acetate/hexane to give title compound as an oil (1.7 g, 80%).


20 D.



- 25 Sodium borohydride (57 mg, 1.51 mmol) was added slowly to a solution of Part C compound (900 mg, 2.1 mmol) in anhydrous methanol (5 mL) at 0°C. The reaction was stirred for 2 hr at 0°C. Additional sodium borohydride (57 mg, 1.51 mmol) was added and stirring was continued for 2 hr and then the reaction mixture was

warmed to room temperature. The mixture was partitioned between ethyl acetate/chloroform and water. The organic layer was washed with water, dried ( $\text{MgSO}_4$ ) and concentrated to give the title compound as a white foam (650 mg, 72%).

E.




10

To a stirred solution of Part D compound (645 mg, 1.5 mmol) in dioxane (8 mL) at room temperature was added zinc chloride (6 mg, 0.045 mmol) followed by thionyl chloride (220  $\mu\text{L}$ , 3.0 mmol). The reaction was stirred for 1 hr and concentrated. The residue was dissolved in ethyl acetate, washed with brine and 5% aqueous sodium bicarbonate solution, dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the title compound as a foam (650 mg, 95%).

20

F.




25

A solution of Part E compound (600 mg, 1.34 mmol) and sodium cyanide (92 mg, 1.88 mmol) in acetonitrile (6 mL) and water (400  $\mu\text{L}$ ) was stirred at reflux overnight. Additional sodium cyanide (92 mg, 1.88 mmol) in water

(400  $\mu$ L) was added and the refluxing continued for 6 hr. Additional sodium cyanide (50 mg, 1.01 mmol) in water (400  $\mu$ L) was added and refluxing was continued for 2 hr. The reaction mixture was cooled to room temperature and concentrated. The residue was partitioned between dichloromethane and water. The organic layer was dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 10:90 to 30:70 ethyl acetate/hexane to give title compound as a white solid (300 mg, 51%).


G.




15

A solution of Part F compound (255 mg, 0.58 mmol) and azidotrimethyltin (167 mg, 0.81 mmol) in xylene (2 mL) was heated at 130°C overnight. The reaction mixture was cooled to room temperature, methanol (8 mL) was added and the mixture was stirred for 30 min. The milky white precipitate was filtered and the filtrate was concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 1:1 ethyl acetate/hexane and then with 95:5 dichloromethane/methanol to give title compound as a white foam (216 mg, 77%).



Example 6


A.



- 5 Nitrogen was bubbled through a solution of Example 5 Part B compound (1.0 g, 4.96 mmol), 3-aminophenyl boronic acid (442 mg, 3.22 mmol) and aqueous sodium carbonate (2 mL, 2M, 4 mmol) in toluene (7 mL) and ethanol (4 mL) at room temperature for 30 min.
- 10 Tetrakis(triphenylphosphine) palladium(0) (95 mg, 0.08 mmol) was added and the mixture was stirred at 80°C overnight under argon. The reaction mixture was cooled to room temperature and extracted with ethyl acetate. The combined organic extracts were washed with brine,
- 15 dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 2:8 to 3:7 ethyl acetate/hexane to give title compound as a white solid (400 mg, 39%).


20

B.



The mixture of Part A compound (400 mg, 0.96 mmol), diisopropylethylamine (333  $\mu$ L, 1.92 mmol), and ethylbromoacetate (95  $\mu$ L, 0.86 mmol) in anhydrous dimethylformamide (3 mL) was stirred at room temperature overnight. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with water, brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 2:8 to 3:7 ethyl acetate/hexane to give title compound as a foam (400 mg, 83%).


15 C.




A solution of Part B compound (400 mg, 0.77 mmol) in aqueous sodium hydroxide (2.3 mL, 1N, 2.3 mmol) and methanol (15 mL) was stirred at room temperature for 20 hr. The reaction mixture was concentrated to a white suspension in water bath at 30°C. The suspension was dissolved in distilled water. The pH was adjusted to 4 by aqueous citric acid (1N). The white precipitate was dissolved in ethyl acetate. The organic layer was washed

with water, brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the title compound as a foam (350 mg, 96%).

5


Example 7

A.




Example 5 Part C compound (214 mg, 0.5 mmol) was added to a solution of silicon tetrachloride (58  $\mu\text{L}$ , 0.5 mmol) and sodium azide (98 mg, 1.5 mmol) in anhydrous acetonitrile (3 mL) at  $0^\circ\text{C}$ . The reaction mixture was warmed to room temperature, then to  $50^\circ\text{C}$  and stirred for 48 hr. The reaction was cooled to room temperature and poured into aqueous saturated sodium carbonate solution. The pH of the mixture was adjusted between 9-10 with sodium carbonate solution and it was extracted with chloroform twice. The combined organic layers were washed with water, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 15:85 to 20:80 ethyl acetate/hexane to give title compound as a foam (56 mg, 26%).

B.




A solution of Part A compound (50 mg, 0.11 mmol) and azidotrimethyltin (34 mg, 0.16 mmol) in xylene (3 mL) was heated at 100°C for 48 hr. The reaction mixture was concentrated to remove all xylene. Fresh xylene (250 µL) and azidotrimethyltin (34 mg, 0.16 mmol) was added and the mixture was heated at 130°C overnight. The reaction mixture was cooled to room temperature, methanol (8 mL) was added, stirred for 30 min, filtered and concentrated. Flash chromatography on silica gel, eluting with a step gradient of 1:4 to 1:1 ethyl acetate/hexane and then with 95:5 to 90:10 dichloromethane/methanol gave the title compound as a foam (45 mg, 87%).

15

Example 8

A.




Nitrogen was bubbled through a solution of Example 5 Part B compound (4.925 g, 17.2 mmol), 3-methoxy phenyl

boronic acid (3.50 g, 23.0 mmol) and 2M aqueous sodium carbonate (14 mL) in toluene (50 mL) and absolute ethanol (20 mL) at room temperature for 30 min.


5 Tetrakis(triphenylphosphine) palladium(0) (795 mg, 0.68 mmol) was added and the mixture was stirred at 80°C for 18 hr under argon. The reaction mixture was cooled to room temperature and partitioned between ethyl acetate/diethyl ether and water. The organic layer was washed with brine, dried over anhydrous MgSO<sub>4</sub> and  
10 concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 1:3 ethyl acetate/hexane to give title compound as a light yellow foam (7.34 g, 99%).

15 B.




To a solution of Part A compound (7.341 g, 17.07 mmol) in anhydrous dichloromethane (100 mL) at 0°C was added boron tribromide in dichloromethane (38 mL, 1M, 38  
20 mmol). The reaction was stirred for 1.5 hr, quenched with aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with water, and brine, then dried over anhydrous MgSO<sub>4</sub>, filtered quickly and concentrated. The solid was  
25 triturated with ether/ethyl acetate/methanol to give the title compound as a white solid (3.156 g, 91% purity). The mother liquor was concentrated and purified by flash chromatography on silica gel, eluting with 15:85 ethyl acetate/hexane to give title compound which was  
30 triturated from ether/dichloromethane to give a white solid (626 mg, >95% purity).

C.



A slurry of Part B compound (91% pure, 3.15 g) in  
5 anhydrous dimethylformamide (35 mL) was warmed to  
partially dissolve the phenol. The mixture was cooled to  
30°C and potassium carbonate (920 mg, 6.6 mmol) was added  
followed by ethyl bromoacetate (1.35 mL, 12.12 mmol).  
The reaction mixture was stirred at room temperature for  
10 24 hr. Additional potassium carbonate (920 mg, 6.6 mmol)  
and ethyl bromoacetate (1.35 mL, 12.12 mmol) was added  
and stirring was continued for 40 hr. The reaction  
mixture was partitioned between ethyl acetate and water.  
The organic layer was washed with water, and brine, then  
15 dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude  
product was purified by flash chromatography on silica  
gel, eluting with 1:9 ethyl acetate/dichloromethane to  
give title compound as a white foam (2.678 g, 77%).


20 D.



A solution of Part C (2.670 g, 5.31 mmol) compound  
in 1,4-dioxane (15 mL) was treated with sodium hydroxide  
solution (13 mL, 1N, 13 mmol). The reaction mixture was  
25 stirred at 50°C for 45 min. The reaction mixture was

cooled to room temperature and acidified with aqueous citric acid solution (25 mL, 10%). The mixture was diluted with water (120 mL) and stirred vigorously for 30 min. The precipitate was filtered, washed with water and  
5 dried to give the title compound as a white solid (2.533 g 100%)

Example 9




10 Neat triethyl phosphonate (201  $\mu$ L, 1.28 mmol) was added dropwise to a stirred suspension of sodium hydride (52 mg, 1.28 mmol) in anhydrous tetrahydrofuran (2 mL). The mixture was stirred at room temperature for 1 hr followed by slow addition of Example 5 Part C compound  
15 (500 mg, 1.16 mmol) in anhydrous tetrahydrofuran (2 mL). The reaction mixture stirred for 5 hr. Additional sodium hydride (20 mg, 0.50 mmol) was added and stirring was continued for 1 hr. The reaction mixture was partitioned between diethyl ether and water. The organic layer was  
20 washed with water, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The residue was purified by flash chromatography on silica gel, eluting with 3:7 ethyl acetate/hexane to give the Example 9 ethyl ester (167 mg, 29%).


25 A portion of the ester was hydrolyzed during the work up. The above aqueous layer was acidified with aqueous citric acid to pH 2. The separated solid was extracted with ether. The organic layer was washed with water, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The  
30 crude product was purified by flash chromatography on

silica gel, eluting with a step gradient of 2:98 to 1:9 methanol/dichloromethane to give title compound as a foam (30 mg, 6%). The title compound is also readily obtained by base hydrolysis of the intermediate ester.

5


Example 10

A solution of Example 5 Part F compound (44 mg, 0.1 mmol) in aqueous sodium hydroxide (250  $\mu$ L, 10N, 2.5 mmol) and ethanol (2 mL) was refluxed for 5 hr and then stirred at room temperature for 14 hr. The reaction mixture was concentrated and the residue was dissolved in water. The pH was adjusted to 4 by acetic acid. The precipitate was filtered and dissolved in chloroform. The organic solution was washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 2:8 ethyl acetate/hexane and then with 95:5 to 90:10 dichloromethane/methanol to give title compound as a foam (34 mg, 62%).

Example 11




A.




A slurry of the ester described in Example 9 (100 mg, 0.20 mmol) and 10% palladium on carbon (20 mg) in methanol (3 mL) was stirred in a parr shaker under hydrogen (40 psi) for 2 hr. The reaction mixture was filtered through a pad of Celite 525 and concentrated to give the title compound as a colorless oil (100 mg, 98%).


10 B.



A solution of Part A compound (100 mg, 0.2 mmol) in aqueous potassium hydroxide (600  $\mu$ L, 2N, 1.2 mmol) and tetrahydrofuran (2 mL) was refluxed for 48 hr. The reaction mixture was concentrated and the residue was dissolved in water. The pH was adjusted to 2 by aqueous HCl (1N). The white precipitate was extracted into ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated to give a foam. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 98:2 to 95:5 dichloromethane/methanol to give title compound as a foam (32 mg, 34%).


Example 12

A.




- 5 Nitrogen was bubbled through a solution of Example 1  
Part A compound (10 g, 26.57 mmol), 1-formyl phenyl  
boronic acid (5.18 g, 34.55 mmol) and aqueous sodium  
carbonate (21.25 mL, 2M, 42.5 mmol) in toluene (70 mL)  
and ethanol (27 mL) at room temperature for 30 min.
- 10 Tetrakis(triphenylphosphine) palladium(0) (1 g, 0.86  
mmol) was added and the mixture was stirred at 80°C for  
14 hr under argon. The reaction mixture concentrated and  
diluted with ethyl acetate/water. The aqueous layer was  
extracted with ethyl acetate (2x200 mL). The combined  
15 organic extracts were washed with brine, dried over  
anhydrous MgSO<sub>4</sub> and concentrated. The crude product was  
purified by flash chromatography on silica gel, eluting  
with a step gradient of 1:19 to 2:8 ethyl acetate/hexane  
to give title compound as a yellow solid (6.11 g, 57.3  
20 %). Less pure fractions were also collected 2.84 g.

B.




Part A compound (1 g, 2.49 mmol) was added to a solution of silicon tetrachloride (285  $\mu$ L, 2.49 mmol) and sodium azide (0.486 g, 7.47 mmol) in anhydrous acetonitrile (10 mL) at 0°C. The reaction mixture was warmed to room temperature, then to 50°C and stirred for 18 hr. The reaction was cooled to room temperature and poured into a mixture of ethyl acetate and aqueous saturated sodium carbonate solution. The pH of the mixture was adjusted between 9-10 with sodium carbonate solution and the aqueous layer was extracted with ethyl acetate (3x100 mL). The combined organic layers were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated the title compound as a yellow oil (1.06 g, 107%).


C.



A solution of Part B compound (0.20 g, 0.50 mmol) and azidotrimethyltin (0.140 g, 0.70 mmol) in p-xylene (2 mL) was refluxed for 18 hr. The reaction mixture was cooled to room temperature, methanol (10 mL) was added and the mixture was stirred for 30 min and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 8:2 ethyl acetate/hexane to 9:1:0.5 dichloromethane/methanol/acetic acid to give title compound as a solid (0.107 g, 48.6 %).

Example 13

A.




5 To a solution of Example 1 Part A compound (7.52 g, 20 mmol) in anhydrous tetrahydrofuran (100 mL) at  $-78^{\circ}\text{C}$  was added dropwise *n*-butyl lithium in hexane (9.3 mL, 2.5 N, 23.25 mmol). The reaction mixture was stirred for 15 min followed by addition of sublimed zinc bromide (5.2 g, 10 23.1 mmol) in anhydrous tetrahydrofuran (40 mL). To this blue solution was added *O*-*t*-butyldimethylsilyl-3-iodophenol (6.7 g, 21.1 mmol) in tetrahydrofuran (40 mL) followed by tetrakis(triphenylphosphine) palladium(0) (1.2 g, 0.96 mmol). The reaction was stirred at  $-78^{\circ}\text{C}$  15 for 30 min and then at room temperature for 20 hr. The reaction mixture was quenched with saturated aqueous ammonium chloride solution and extracted with diethyl ether thrice. The combined organic layers were dried over anhydrous  $\text{MgSO}_4$  and concentrated to get crude amber 20 oil.

To a solution of the crude amber oil in anhydrous tetrahydrofuran (100 mL) was added dropwise tetrabutyl ammonium fluoride in tetrahydrofuran (25 mL, 1 M, 25 mmol). The solution was stirred at room temperature for 25 1 hr, quenched with saturated aqueous ammonium chloride solution and extracted with diethyl ether thrice. The

combined organic layers were washed with saturated aqueous ammonium chloride solution, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 2:8 ethyl acetate/hexane to give title compound as a yellow foam (4.6 g, 59%).


B.




To a slurry of Part A compound (200 mg, 0.51 mmol) and cesium carbonate (217 mg, 0.67 mmol) in anhydrous dimethylformamide (10 mL) was added ethyl 2-bromopropionate (121 mg, 0.67 mmol). The reaction mixture was stirred at room temperature for 18 h, diluted with water (20 mL), extracted with ethyl acetate (3x20 mL). The organic layer was washed with water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated to get the crude ester as a yellow oil.

A solution of crude ester in aqueous sodium hydroxide solution (5 mL, 1N) and dioxane (5 mL) was refluxed for 20 min. The reaction mixture was cooled, concentrated, diluted with water and pH was adjusted to 1 with HCl (1N). The mixture was extracted with ethyl acetate (3x20 mL). The combined organic layers were washed with water, and brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 75:25 ethyl acetate/methanol to give title compound as a off-white powder (208 mg, 88%).

30

Example 14


A solution of Example 1 Part E compound (165 mg, 0.4 mmol) in aqueous sodium hydroxide (1 mL, 10N, 10 mmol) and ethanol (3 mL) was refluxed for 5 h and then stirred at room temperature for 14 h. The reaction mixture was concentrated and the residue was dissolved in water. The pH was adjusted to 4 by acetic acid. The precipitate was filtered and dissolved in chloroform. The organic solution was filtered and the filtrate was concentrated to get the title compound as an off-white solid (95 mg, 55%).

Example 15

To a solution of Example 1 Part C compound (75 mg, 0.18 mmol) in tetrahydrofuran (4 mL) at 0°C was added sulfamic acid (19 mg, 0.18 mmol) as a solid. To this cold solution was added dropwise a solution of sodium chlorite (20 mg, 0.18 mmol) in water (2 mL) over 10 min. The reaction was stirred for 1 hr and then partitioned between ether and brine. The organic layer was dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient 2:8 ethyl acetate/hexane to


5:95 methanol/dichloromethane to give title compound as a white solid (53 mg, 71%).

Example 16




5

A.




To a solution of Example 13 Part A compound (389 mg, 1 mmol) and triethylamine (420  $\mu$ L, 3 mmol) in  
10 dichloromethane (25 mL) at 0°C was added slowly triflic anhydride (185  $\mu$ L, 1.1 mmol). The reaction was warmed to room temperature over 2 hr, washed with water, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was  
15 purified by flash chromatography on silica gel, eluting with 1:4 ethyl acetate/hexane to give title compound as a brown oil (485 mg, 93%).

B.



A mixture of Part A compound (485 mg, 0.93 mmol), triethylamine (142  $\mu$ L, 1.02 mmol), 1,3-bis(diphenylphosphino) propane (11 mg, 0.02 mmol), methyl acrylate (167  $\mu$ L, 1.86 mmol) and palladium acetate (5.2 mg, 0.02 mmol) in dimethylformamide (10 mL) was refluxed overnight at 100°C. The reaction mixture was cooled to room temperature, diluted with dichloromethane. The organic layer was washed with aqueous 1 N HCl, water, brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 15:85 ethyl acetate/hexane to give title compound as an off-white powder (234 mg, 55%).


C.




15

A solution of Part B compound (234 mg, 0.51 mmol) in 1,4-dioxane (5 mL) and sodium hydroxide (5 mL, 1M, 5 mmol) was refluxed for 0.5 hr. The reaction mixture was cooled to room temperature, pH was adjusted to 2 with HCl solution (1 M), and then extracted with ethyl acetate (2x20 mL). The combined organic layers were washed water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the title compound as a white solid (198 mg, 87%). mp 190-192°C.




Example 17

A slurry of Example 16 Part C compound (136 mg, 0.31 mmol) and 10% palladium on carbon (100 mg) in  
5 tetrahydrofuran (20 mL) was stirred under hydrogen atmosphere overnight. The reaction was filtered and concentrated. The residue was dissolved in ethyl acetate, washed with water and brine, dried over anhydrous MgSO<sub>4</sub> and concentrated to give the title  
10 compound as a white solid (124 mg, 90%). mp 91-93°C.

Example 18

A.




15

1-Methyl-3-nitro-1-nitrosoguanidine was added to a mixture of ether and aqueous potassium hydroxide solution at 0°C. The yellow ether layer was decanted into a flask containing solid potassium hydroxide cooled to 0°C. The  
20 ether was decanted into a solution of Example 16 Part B compound in dichloromethane at 0°C. Palladium acetate


was added to this mixture and the reaction was stirred for 12 h. The reaction mixture was filtered through a pad of Celite 525 and the filtrate was concentrated to give a brown oil. The crude product was purified by  
5 flash chromatography on silica gel, eluting with 1:4 ethyl acetate/hexane to give title compound as a yellow oil (90 mg g, 92%).

B.




10

A solution of Part A (90 mg, 0.203 mmol) and lithium hydroxide (9 mg, 0.223 mmol) in tetrahydrofuran (5 mL) and water (5 mL) was stirred at room temperature overnight. Acidified with concentrated HCl to pH < 7,  
15 extracted with ethyl acetate (3X20mL). The combined organic layers were washed with water, and brine, then dried over MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 4:6 ethyl acetate/hexane to give title compound as a  
20 white foam (82 mg, 88%).

Example 19


A.



5

To a solution of 2'-bromoacetophenone (6.7 mL, 50 mmol) and potassium hydroxide (14.03 g, 150 mmol) in methanol (100 mL) was added benzaldehyde (15.3 mL, 150 mmol) at room temperature. The reaction mixture was stirred for 30 min, and poured into a mixture of acetic acid and water (1:2, 300 mL) at 0°C. The mixture was extracted with dichloromethane (3x100 mL). The combined organic layers were washed with saturated aqueous ammonium chloride solution, brine, then dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was distilled under vacuum to remove benzaldehyde. The residue was purified by flash chromatography eluting with 9:11 dichloromethane/hexane to give title compound as a light yellow oil (10.4 g, 72%).

B.




25

A solution of Part A compound (9.18 g, 31.9 mmol), benzaldehyde (3.4 mL, 33.5 mmol), triethylamine (2.6 mL,

18.5 mmol) and 3,4-dimethyl-5-(2-hydroxyethyl)thiazolium iodide (1.82 g, 6.4 mmol) in ethanol (45 mL) was refluxed for 40 hr. The orange red solution was evaporated under reduced pressure to remove ethanol. The residue was  
5 dissolved in dichloromethane (200 mL). The organic layer was washed with dilute HCl (1M, 200 mL), saturated aqueous sodium bicarbonate solution (200 mL), water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The orange red residue was purified by flash  
10 chromatography eluting with a step gradient of 50% to 66% dichloromethane in hexane to give title compound as a colorless oil (9.80 g, 78%).


C.



15


A solution of Part B compound (1.31 g, 3.34 mmol) and boron trifluoride diethyl etherate (0.42 mL, 3.34  
20 mmol) in toluene (15 mL) was refluxed for 3 hr. The reaction mixture was cooled to room temperature and diluted with hexane (250 mL). The organic layer was washed with saturated aqueous sodium bicarbonate solution (200 mL), water, and brine, then dried over anhydrous  
25  $\text{MgSO}_4$  and concentrated. The light yellow oil was purified by flash chromatography eluting 1:9 dichloromethane/hexane to give title compound as a colorless oil (1.10 g, 88%).

D.



5 Nitrogen was bubbled through a solution of Part C compound (1.0 g, 2.66 mmol), 3-methoxy phenylboronic acid (526 mg, 3.46 mmol) and sodium carbonate (9.3 mL, 2M, 18.6 mmol) in toluene (7 mL) and ethanol (2.8 mL) at room temperature for 30 min. Tetrakis(triphenylphosphine)-  
10 palladium(0) (100 mg, 0.086 mmol) was added and the reaction mixture was stirred at 90°C for 16 h under nitrogen. The reaction mixture was cooled to room temperature and extracted with ethyl acetate (2x50 mL). The combined organic layers were washed with brine, dried  
15 over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography eluting with 1:3 dichloromethane/hexane to give title compound as a white foam (1.01 g, 94%).


20 E.



To a mixture of Part D compound (791 mg, 1.97 mmol)  
25 and lithium iodide (3.95 g, 29.5 mmol) was added dimethyl formamide (7 mL) at room temperature. The reaction mixture was refluxed at 170°C overnight under argon, cooled to room temperature, diluted with water and extracted with ether (3x35 mL). The combined organic  
30 layers were washed with saturated aqueous sodium

bicarbonate solution (80 mL), water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 1:3 ethyl acetate/hexane to give title compound as a white foam (651 mg, 85%).

F.




10

To a solution of Part E compound (368 mg, 0.95 mmol) in anhydrous dimethylformamide (2.5 mL) was added sodium hydride (38 mg, 60%, 0.95 mmol). The reaction was stirred at room temperature for 20 min followed by addition of bromoacetonitrile (0.10 mL, 1.44 mmol). The reaction mixture was stirred at 50°C overnight, cooled to room temperature, diluted with water and extracted with ether (3x40 mL). The combined organic layers were washed with saturated aqueous sodium bicarbonate solution (80 mL), water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 1:9 dichloromethane/hexane to give title compound as a white foam (306 mg, 75%).


25

G.




A solution of Part F compound (174 mg, 0.41 mmol) and azidotrimethyltin (126 mg, 0.61 mmol) in toluene (2 mL) was refluxed overnight. The reaction mixture was cooled to room temperature, methanol (5 mL) was added, stirred for 30 min, and concentrated. The residue was dissolved in dichloromethane (100 mL), washed with water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 1:9 methanol/ethyl acetate to give title compound as a white solid (110 mg, 57%).

#### Example 20




15           A.




A solution of Example 19 Part B compound (787 mg, 2.0 mmol) and ammonium acetate (800 mg, 10.4 mmol) in acetic acid (5 mL) was refluxed, protected by a drying tube, for 3 hr. The reaction mixture was cooled to room temperature and poured slowly into aqueous saturated sodium bicarbonate and the reaction mixture was extracted with ethyl acetate. The organic layer was dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 11:39 dichloromethane/hexane, to give title compound as a white foam (508 mg, 68%).

B.



Nitrogen was bubbled through a solution of Part A  
5 compound (775 mg, 2.07 mmol), 3-formyl phenylboronic acid  
(405 mg, 2.7 mmol) and aqueous sodium carbonate (3.3 mL,  
2 M, 6.6 mmol) in toluene (5.5 mL) and ethanol (2.2 mL)  
at room temperature for 30 min.  
Tetrakis(triphenylphosphine)-palladium(0) (80 mg, 0.07  
10 mmol) was added and the reaction mixture was stirred at  
80°C for 14 hr under argon. The reaction mixture was  
cooled to room temperature and extracted with ethyl  
acetate twice. The combined organic extracts were washed  
with brine, dried (MgSO<sub>4</sub>) and concentrated. The crude  
15 product was purified by flash chromatography on silica  
gel, eluting with 2:1 dichloromethane/hexane to give  
title compound as a white foam (598 mg, 72%).

C.




20

Lithium borohydride in tetrahydrofuran (0.5 mL, 2M,  
1 mmol) was added slowly to a solution of Part B compound  
(495 mg, 1.24 mmol) in tetrahydrofuran (3 mL) at room  
temperature under argon. The reaction was stirred for 30  
25 min, quenched cautiously with aqueous saturated sodium  
bicarbonate solution, stirred for 10 min and extracted  
with ethyl acetate. The organic extract was dried




(MgSO<sub>4</sub>) and concentrated. The residue was dissolved in dichloromethane and filtered through a pad of silica gel (25 g). The pad was washed with dichloromethane (200 mL). The filtrate was dried (MgSO<sub>4</sub>) and concentrated to  
5 give the title compound as a white foam (490 mg, 98%).

D.




To a solution of Part C compound (214 mg, 0.53 mmol)  
10 and carbon tetrabromide (200 mg, 0.60 mmol) in dichloromethane (2 mL) was added triphenylphosphine (158 mg, 0.53 mmol) at room temperature. The solution was stirred for 1 hr, evaporated onto silica gel and purified by flash chromatography on silica gel, eluting with 3:7  
15 dichloromethane/hexane to give the title compound (180 mg, 73%) as a light yellow foam.

E.




A solution of Part D compound (175 mg, 0.37 mmol)  
20 and sodium cyanide (200 mg, 0.60 mmol) in dimethylsulfoxide (1 mL) was stirred for 3 hr at 50°C under argon. The reaction mixture was cooled to room temperature, diluted with water and filtered. The solid  
25 was washed with water and dried to give the title compound as an off-white solid (140 mg, 91%).

F.




A solution of Part E compound (135 mg, 0.33 mmol)  
5 and azidotrimethyltin (100 mg, 0.49 mmol) in toluene (2 mL) was heated to 140°C for 18 hr under argon. The reaction mixture was cooled to room temperature. After adding methanol (5 mL), the reaction was stirred for 30 min and concentrated. The crude product was purified by  
10 flash chromatography on silica gel, eluting with 1:5 methanol/dichloromethane to give title compound as a tan solid (105 mg, 70%).

Example 21

15


A.



A solution of Example 19 Part B compound (1.164 g, 2.96 mmol), ethylamine hydrochloride (1.22 g, 15 mmol)  
20 and sodium acetate (1.23 g, 15 mmol) in acetic acid (10 mL) was refluxed for 52 h under argon. The reaction


mixture was cooled to room temperature and added slowly to saturated aqueous sodium bicarbonate solution (100 mL). The mixture was extracted with dichloromethane (2 x 100 mL) and the combined organic layers were washed, dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography eluting with 1:7 dichloromethane/hexanes to give title compound as a white foam (556 mg, 47%).

10 B.




Nitrogen was bubbled through a solution of Part A compound (547 mg, 1.35 mmol), 3-amino phenylboronic acid (327 mg, 2.4 mmol) and aqueous sodium carbonate (3.0 mL, 2M, 6 mmol) in toluene (5 mL) and ethanol (2 mL) at room temperature for 30 min. Tetrakis(triphenylphosphine)-palladium(0) (55 mg, 0.05 mmol) was added and the reaction mixture was stirred at 80°C for 14 h under argon. The reaction mixture was cooled to room temperature and extracted twice with ethyl acetate. The combined organic layers were washed with brine, dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography eluting with 11:9 dichloromethane/hexanes to give title compound as a white foam (379 mg, 68%).

C.



To a slurry of Part B compound (373 mg, 0.90 mmol) and potassium carbonate (200 mg, 1.4 mmol) in anhydrous DMF (2 mL) was added bromoacetonitrile (70  $\mu$ L, 1.0 mmol). The reaction mixture was stirred at 50°C overnight, cooled to room temperature, diluted with water, and extracted with ether (3x50 mL). The combined organic layers were washed with water and brine, then dried (MgSO<sub>4</sub>) and concentrated. The crude product was purified by flash chromatography eluting with 3:7 dichloromethane/hexanes to give title compound as a colorless oil (174 mg, 43%).


15 D.



A solution of Part C compound (120 mg, 0.38 mmol) and azidotrimethyltin (100 mg, 0.5 mmol) in toluene (2 mL) was refluxed overnight under argon. The reaction mixture was cooled to room temperature, methanol (5 mL) was added, stirred for 1 hr, and concentrated to give crude title compound as a yellow oil (140 mg). A portion of the impure product (85 mg) was purified by preparative HPLC (C-18 reverse phase, eluting with methanol/0.1%


aqueous TFA) to give the title compound (47 mg, 34%) as a white solid, mp 109-111°C.

Example 22




5

A.



A solution of Example 19 Part B compound (3.26 g, 8.3 mmol) and Lawesson's reagent (7 g, 17 mmol) in toluene (15 mL) was refluxed for 24 hr under argon. Additional Lawesson's reagent (2 g, 4.94 mmol) was added and the solution was refluxed for an additional 24 hr. The reaction mixture was cooled to room temperature and diluted with ether. The organic layer was washed with saturated aqueous sodium bicarbonate solution, water and brine, then dried (MgSO<sub>4</sub>) and concentrated. The crude product was purified by flash chromatography, eluting with 1:24 dichloromethane/hexanes to give title compound as a white solid (2.65 g) with the corresponding furan as an impurity (20%). The mixture was recrystallized from hexanes to give the product as a white solid (850 mg, 93% pure, 28% yield).


B.



Nitrogen was bubbled through a solution of Part A compound (663 mg, 1.69 mmol), 3-aminophenylboronic acid (410 mg, 2.20 mmol) and aqueous sodium carbonate (3.7 mL, 2M, 7.4 mmol) in toluene (6 mL) and ethanol (2.5 mL) at room temperature for 30 min.

Tetrakis(triphenylphosphine)-palladium(0) (67 mg, 0.06 mmol) was added and the reaction mixture was refluxed for 16 h under argon. The reaction mixture was cooled to room temperature, diluted with brine (10 mL) and extracted with ethyl acetate. The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The crude product was purified by flash chromatography, eluting with 3:1 dichloromethane/hexanes to give title compound as a white foam (680 mg, 100%).

C.




To a slurry of Part B compound (354 mg, 0.878 mmol) and potassium carbonate (250 mg, 1.8 mmol) in anhydrous DMF (3 mL) was added *tert*-butyl bromoacetate (130  $\mu$ L, 0.88 mmol). The reaction mixture was stirred at 50°C under argon overnight, cooled to room temperature, diluted with water and extracted with ether (3 x 20 mL). The combined organic layers were washed with water and

brine, then dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography eluting with 2:3 dichloromethane/hexanes to give title compound as a colorless oil (118 mg, 26%).


5

D.




A solution of Part C compound (115 mg, 0.222 mmol) in trifluoroacetic acid (1 mL) and dichloromethane (1 mL) was stirred at room temperature for 14 h under argon. The reaction mixture was evaporated from ethanol (4 mL) twice. The residue was dissolved in ethyl acetate and hexane was added to give an oily product. The solvent was decanted and the product was dried under vacuum at 60°C to give the title compound as a white foam (47 mg, 62%), 87% pure and containing 7.8% of the analogous furan.

20


Example 23

A.



Nitrogen was bubbled through a solution of Example 19 Part C compound (1.0 g, 2.66 mmol), 3-formyl  
5 phenylboronic\_acid (520 mg, 3.5 mmol) and aqueous sodium carbonate (9.3 mL, 2M, 18.6 mmol) in toluene (7 mL) and ethanol (2.8 mL) at room temperature for 30 min. Tetrakis(triphenyl-phosphine) palladium(0) (100 mg, 0.09 mmol) was added and the mixture was stirred at 80°C for  
10 14 hr under argon. The reaction mixture was cooled to room temperature and extracted with ethyl acetate (2x30 mL). The combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The crude product  
15 was purified by flash chromatography on silica gel, eluting with 1:3 dichloromethane/hexane to give title compound as a white foam (525 mg, 49%).

B.




20 Lithium aluminum hydride in tetrahydrofuran (0.7 mL, 1M, 0.7 mmol) was added slowly to a solution of Part A compound (494 mg, 1.23 mmol) in tetrahydrofuran (3 mL) at room temperature under argon. The reaction was stirred for 15 min, quenched cautiously with aqueous potassium  
25 hydrogen sulfate solution (5%) and extracted with ethyl acetate. The organic layer was dried (MgSO<sub>4</sub>) and



concentrated to give the title compound as a white foam (471 mg, 95%).

C.




5

To a solution of Part B compound (465 mg, 1.15 mmol), triphenylphosphine (303 mg, 1.16 mmol) and imidazole (172 mg, 2.5 mmol) in tetrahydrofuran (5 mL) under argon was added a solution of iodine (293 mg, 1.15 mmol) in THF (1 mL) over 5 min. The reaction mixture was stirred for 10 min and then diluted with ethyl acetate. The organic layer was washed with sodium bisulfite solution (5%), dried ( $\text{MgSO}_4$ ) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with hexane to give title compound as a colorless oil (304 mg, 51%).

15

D.




20


A solution of Part C compound (299 mg, 0.584 mmol) and potassium cyanide (300 mg, 4.6 mmol) in DMSO (3 mL) was stirred at 50°C for 13 h. The reaction mixture was cooled to room temperature, diluted with water, and extracted with diethyl ether (3x20 mL). The organic layer was dried ( $\text{MgSO}_4$ ) and concentrated to give the title compound as a yellow oil (193 mg, 80%).

25


E.



A solution of Part D compound (190 mg, 0.46 mmol) and azidotrimethyltin (140 mg, 0.68 mmol) in toluene (3 mL) was refluxed for 18 h under argon. The reaction mixture was cooled to room temperature, methanol (5 mL) was added and the mixture was stirred for 30 min. After evaporating the solvents, the crude product was recrystallized from ethyl acetate/hexane to give title compound as a white solid (77 mg, 37%), mp 188-190°C.

Example 24

A.




15

To a solution of 2-bromo-3-methyl benzoic acid (2.15 g, 10 mmol), 4-dimethyl amino pyridine (211 mg, 1.73 mmol) and benzoin (2.07 g, 9.56 mmol) in dichloromethane (30 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.09 g, 10.6 mmol) at room temperature. The resulting pale yellow solution was

stirred for 48 hr, treated with HCl (20 mL, 0.5 N) and extracted with ethyl acetate (2x150 mL). The combined organic layers were washed with HCl (20 mL, 0.5 N), saturated sodium bicarbonate solution (20 mL), and brine (20 mL), dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 5:95 ethyl acetate/hexane to give the ketoester as a colorless oil (2.83 g, 67%).

A mixture of crude ketoester (2.81 g, 6.9 mmol) and ammonium acetate (2.63 g, mmol) in glacial acetic acid (25 mL) was stirred at 110°C for 4 h. The reaction mixture was concentrated to half the volume, cooled to 0°C and diluted with water. The mixture was extracted with ethyl acetate (2x150 mL). The combined organic layers were washed with water (3x30 mL), saturated aqueous sodium bicarbonate solution (2x30 mL), and brine (30 mL), dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 5:95 ethyl acetate/hexane to give title compound as a white solid (2.46 g, 91%). mp 83-85°C.


B.



To a solution of Part A compound (500 mg, 1.28 mmol) in anhydrous tetrahydrofuran (7 mL) at -78°C was added dropwise n-butyl lithium in hexane (600 µL, 2.5 N, 1.47 mmol). The reaction mixture was stirred for 15 min followed by addition of zinc bromide (330 mg, 1.47 mmol) in anhydrous tetrahydrofuran (3.3 mL). To this light

green solution was added 1-(t-butyldimethylsilyloxy)-3-iodobenzene (428 mg, 1.37 mmol) in tetrahydrofuran (1 mL) followed by tetrakis(triphenylphosphine) palladium(0) (74 mg, 0.06 mmol). The reaction was stirred at -78°C for 5 30 min and then at room temperature for 20 h. The reaction was quenched with aqueous ammonium chloride solution (5 mL, 25%) and extracted with diethyl ether (2x25 mL). The combined organic layers were washed with aqueous ammonium chloride solution (5 mL, 25%), brine (5 10 mL), dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting 2:98 ethyl acetate/hexane to give title compound as a syrup (229 mg, 34.6%).


15 C.



To a solution of Part B compound (229 mg, 0.44 mmol) in anhydrous tetrahydrofuran (4 mL) was added dropwise tetrabutylammonium fluoride in tetrahydrofuran (400  $\mu\text{L}$ , 1 20 M, 0.4 mmol). The yellow solution was stirred at room temperature for 72 h, quenched with aqueous ammonium chloride solution (2 mL, 25%) and extracted with diethyl ether (2x25 mL). The combined organic layers were washed with aqueous ammonium chloride solution (2 mL, 25%), 25 dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 2:98 to 5:95 ethyl acetate/hexane to give title compound as a syrup (163 mg, 92%).


30

D.




To a slurry of Part C compound (41 mg, 0.10 mmol) and potassium carbonate (25 mg, 0.18 mmol) in anhydrous dimethylformamide (0.5 mL) was added ethyl bromoacetate (15  $\mu$ L, 0.14 mmol). The reaction mixture was stirred at room temperature for 20 h, diluted with ethyl acetate (20 mL). The organic layer was washed with water (3x1 mL), and brine (1 mL), then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude ester was used directly for the next step.


E.



A solution of Part D (41 mg, 0.084 mmol) compound and sodium hydroxide solution (250  $\mu$ L, 1N, 0.25 mmol) in methanol (1 mL) and water (1 mL) was stirred at room temperature for 1.5 h. The reaction mixture was concentrated, diluted with water and HCl (200  $\mu$ L, 1N), extracted with ethyl acetate (2x10 mL). The combined organic layers were washed with water (2x1 mL), and brine (1 mL), dried over anhydrous  $\text{MgSO}_4$  and concentrated to give the title compound as a beige foam (41 mg, 100%).

Example 25


A.



5        A solution of Example 1 Part A compound (1.25 g, 3.22 mmol) and tetrakis(triphenylphosphine) palladium(0) (117 mg, 0.10 mmol) in toluene (7.5 mL) was stirred for 10 min. 3-Aminophenyl boronic acid (637 mg, 3.96 mmol) was added to the solution followed by aqueous sodium carbonate (3.3 mL, 2M, 6.6 mmol). The reaction mixture was stirred at 80°C for 20 hr under argon, cooled to room temperature and concentrated. The residue was partitioned between ethyl acetate (2x90 mL) and water (9 mL). The combined organic layers were washed with brine, 15 dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 1:9 to 1:4 ethyl acetate/hexane to give title compound as an oil (1.445 g, 100%).


20

B.



The mixture of Part A compound (1.445 g, 3.22 mmol), diisopropylethylamine (1.12 mL, 6.44 mmol), and  
5 benzylbromoacetate (610  $\mu$ L, 3.7 mmol) in anhydrous dimethylformamide (15 mL) was stirred at room temperature for 17 hr. The reaction mixture was diluted with ethyl acetate (250 mL). The organic layer was washed with  
10 water (3x40 mL), brine (40 mL), dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 1:9 to 1:4 ethyl acetate/hexane to give title compound as an oil (1.57 g, 91%).


15 C.



To a slurry of Part B compound (100 mg, 0.19 mmol) and potassium carbonate (26 mg, 0.19 mmol) in anhydrous dimethylformamide (1 mL) was added methyl iodide (17  $\mu$ L,  
20 0.28 mmol). The reaction mixture was stirred at room temperature for 16 h. Additional methyl iodide (60  $\mu$ L, 0.99 mmol) was added and the reaction was stirred for 24 hr. Methyl iodide (40  $\mu$ L, 0.66 mmol) was added again and the reaction was stirred for 20 hr. The reaction mixture


was diluted with ethyl acetate (25 mL). The organic layer was washed with water (3x2 mL), and brine (2 mL), then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The golden oil was purified by flash chromatography on silica  
5 gel, eluting with 1:9 ethyl acetate/hexane to give title compound as colorless oil (73 mg, 70%).

D.




10 A suspension of Part C compound (73.5 mg, 0.13 mmol) and 20% palladium hydroxide on carbon (12.9 mg) in ethyl acetate (3 mL) was stirred under hydrogen atmosphere (1 atm) for 5 h. Additional catalyst (12 mg) was added and the reaction was stirred for 5 h. The reaction mixture  
15 was filtered through a pad of Celite® 525. The solids were washed with ethyl acetate (2x10 mL) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 1:4 ethyl acetate/hexane to 95:5  
20 dichloromethane/methanol to give title compound as a foam (14 mg, 24%).

#### Example 26






A.



A solution of Example 24 Part A compound (700 mg, 1.79 mmol) and tetrakis(triphenylphosphine) palladium(0) (63 mg, 0.055 mmol) in toluene (4 mL) was stirred for 10 min. 3-aminophenyl boronic acid (344 mg, 2.15 mmol) was added to the solution followed by aqueous sodium carbonate (1.8 mL, 2M, 3.6 mmol). The reaction mixture was stirred at 80°C for 20 h under argon, cooled to room temperature and concentrated. The residue was partitioned between dichloromethane (2x50 mL) and water (5 mL). The combined organic layers were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 1:9 ethyl acetate/hexane to give title compound as a white foam (623 mg, 86%).

B.




The mixture of Part A compound (300 mg, 0.72 mmol), diisopropylethylamine (0.25 mL, 1.44 mmol), and benzylbromoacetate (130 µL, 0.79 mmol) in anhydrous dimethylformamide (3 mL) was stirred at room temperature for 20 hr. The reaction mixture was diluted with ethyl acetate (100 mL). The organic layer was washed with water (2x10 mL), brine (10 mL), dried over anhydrous

MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 5:95 to 1:9 ethyl acetate/hexane to give title compound as a white foam (341 mg, 86%).


5

C.




A suspension of Part B compound (311 mg, 0.56 mmol) and 20% palladium hydroxide on carbon (50 mg) in ethyl acetate (10 mL) was stirred under hydrogen atmosphere (1 atm) for 5 h. The reaction mixture was filtered through a pad of Celite® 525. The solids were washed with ethyl acetate (3x10 mL) and concentrated. The crude product was triturated with 1:5 dichloromethane/pentane (25 mL) to give the title compound as an off-white precipitate which was washed with pentane and dried (222 mg, 86%). mp 175-177°C.

#### Example 27



20


A.



To a solution of 2-(3-nitro-p-toluy1)-benzoic acid (325 mg, 1.26 mmol), 4-dimethylaminopyridine (41 mg, 0.34 mmol) and benzoin (404 mg, 1.90 mmol) in dichloromethane (6 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (408 mg, 2.09 mmol) at room temperature. The resulting pale yellow solution was stirred for 16 h, treated with HCl (4 mL, 0.5 N) and extracted with ethyl acetate (2x30 mL). The combined organic layers were washed with HCl (4 mL, 0.5 N), saturated sodium bicarbonate solution (4 mL), and brine (4 mL), dried over anhydrous MgSO<sub>4</sub> and concentrated to give a light yellow oil.


A mixture of crude ketoester and ammonium acetate (500 mg, 6.49 mmol) in glacial acetic acid (5 mL) was stirred at 105°C for 4 h. The reaction mixture was concentrated and extracted with ethyl acetate (2x50 mL). The combined organic layers were washed with water (3x12 mL), saturated aqueous sodium bicarbonate solution (2x12 mL), and brine (7 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 5:95 ethyl acetate/hexane to give title compound as a white foam (265 mg, 48%). mp 108-110°C.

B.



A suspension of Part A compound (265 mg, 0.61 mmol) and 20% palladium hydroxide on carbon (50 mg) in ethyl acetate (10 mL) was stirred under hydrogen atmosphere (1 atm) for 41 h. The reaction mixture was filtered through a pad of Celite® 525. The solids were washed with ethyl acetate (3x10 mL) and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 5:95 to 1:9 ethyl acetate/hexane to give title compound as a white foam (259 mg, 100%).

C.




15

The mixture of Part B compound (255 mg, 0.63 mmol), diisopropylethylamine (0.22 mL, 1.26 mmol), and benzylbromoacetate (120 µL, 0.72 mmol) in anhydrous dimethylformamide (3 mL) was stirred at room temperature for 20 h. The reaction mixture was diluted with ethyl acetate (250 mL). The organic layer was washed with water (3x8 mL), brine (8 mL), dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 1:9

ethyl acetate/hexane to give title compound as a white foam (226 mg, 65%).

D.




5


A suspension of Part C compound (226 mg, 0.41 mmol) and 20% palladium hydroxide on carbon (37 mg) in ethyl acetate (7.5 mL) was stirred under hydrogen atmosphere (1 atm) for 5 h. The reaction mixture was filtered through a pad of Celite® 525. The solids were washed with ethyl acetate (3x10 mL) and concentrated. The crude product was triturated with 1:4 dichloromethane/hexane (25 mL) to get the title compound as a cream colored precipitate which was washed with hexane and dried (188 mg, 100%). mp 182-184°C.

15

Example 28




A.



20


2-Bromobenzoyl chloride (15.2 mL, 0.118 mol) was added dropwise to a stirred solution of phenyl hydrazine (11.6 mL, 0.118 mol) and triethylamine (16.0 mL, 0.115 mol) in ether (400 mL) at 0°C over 30 min. The reaction  
5 was warmed to room temperature. The solids formed were filtered and washed with ether thrice. The solids were dissolved in dichloromethane washed with water, and brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude was recrystallized from ethyl acetate to get the  
10 title compound (12.1 gm, 35%)

B.




A solution of Part A compound (5 gm, 17.2 mmol) and  
15 phosphorus pentachloride (4.22 gm, 19.3 mmol) in ether (100 mL) was refluxed for 19 hr. The reaction mixture was cooled to room temperature and treated with a solution of phenol (7.38 gm, 78.4 mmol) in ether (10 mL). After 10 min, methanol (7.3 mL) was added, the mixture  
20 was stirred for 5 min and concentrated. The residue was purified by flash chromatography on silica gel eluting with 10% EtOAc in hexane to give title compound as a light brown 0:1 (2.07 g, 39%).

25 C.



A solution of Part B compound (2.0 gm, 6.46 mmol) in absolute ethanol (9.4 mL) was added to a solution of ethylbenzoyl acetate (1.26 mL, 6.55 mmol) and 21% sodium ethoxide in ethanol (2.09 mL, 6.44 mmol) in absolute ethanol (12.6 mL). The reaction was stirred at room temperature for 4 h, quenched with 2N HCl (7.5 mL) and concentrated. The residue was triturated with ether. The organic layer was washed with water, and brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The residue was purified by flash chromatography on silica gel eluting with a step gradient of 5% to 10% EtOAc in hexane to give the title compound as a thick oil (1.1 g, 38%).

D.




15

Nitrogen was bubbled through a solution of Part C compound (1.145 g, 2.56 mmol), 3-methoxyphenyl boronic acid (478 mg, 3.15 mmol) and aqueous sodium carbonate (2.6 mL, 2M, 5.2 mmol) in toluene (6 mL) and ethanol (2.6 mL) at room temperature for 15 min.


Tetrakis(triphenylphosphine) palladium(0) (93 mg, 0.08 mmol) was added and the mixture was stirred at 80°C for 20 h under argon. The reaction mixture was cooled to room temperature, diluted with water and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 5% to 10% ethyl acetate in hexane to give the title compound (1.215 g, 100%).

E.



To a solution of Part D compound (948 mg, 1.99 mmol) in dry dichloromethane (10 mL) at 0°C was added boron tribromide in dichloromethane (1M, 4.1 mL, 4.1 mmol). The reaction was stirred at 0°C for 3 h and then at room temperature overnight. The mixture was diluted with water followed by 1N HCl and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 10% to 50% ethyl acetate in hexane to give title compound (688 mg, 75%).

F.




A solution of Part E compound (670 mg, 1.45 mmol) in anhydrous ether (3.6 mL) was added to a slurry of lithium aluminium hydride (114 mg, 2.85 mmol) in anhydrous ether (6 mL) at 0°C. The reaction was stirred at 0°C for 3 h and at room temperature for 4 h. Anhydrous THF (10 mL) was added to the reaction mixture and it was stirred for 4 h. The reaction was cooled to 0°C and quenched with 10% HCl and extracted with ethyl acetate. The combined



organic extracts were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with a step gradient of 10% to 20% to 50% ethyl acetate  
5 in hexane to give the title compound (474 mg, 78%).


G.



To a slurry of Part F compound (472 mg, 1.13 mmol)  
10 and potassium carbonate (180 mg, 1.3 mmol) in anhydrous dimethylformamide (5 mL) was added ethyl bromoacetate (150  $\mu\text{L}$ , 1.3 mmol). The reaction mixture was stirred at room temperature for 48 h, diluted with ethyl acetate and water. The organic layer was washed with water and  
15 brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude was purified by flash chromatography on silica gel, eluting with a step gradient of 25% to 33% ethyl acetate in hexane to give the title compound as an oil (410 mg, 74%).


20

H.




A solution of Part G compound (60 mg, 0.12 mmol) in aqueous sodium hydroxide (350  $\mu\text{L}$ , 1N, 3.5 mmol) and  
25 methanol (1.4 mL) was heated to 50°C for 3 h. The


reaction mixture was concentrated and the residue was diluted with water. The pH was adjusted to 1 with 1N HCl. The mixture was extracted with ethyl acetate. The combined organic layers were washed with brine, dried  
5 over anhydrous  $\text{MgSO}_4$  and concentrated to give title compound as a foam (63.8 mg, 100%).

Example 29

10 A.



A solution of




generated in analogy to the procedure used in Example 28  
15 Part B, (2 g, 6.46 mmol) in anhydrous benzene (10 mL) was added to a solution of aniline (650  $\mu\text{L}$ , 7.13 mmol) and triethyl amine (970  $\mu\text{L}$ , 6.96 mmol) in anhydrous benzene (10.8 mL) at 50°C. The reaction was stirred at 50°C for 3 hrs and concentrated. The residue was partitioned  
20 between water and ethyl acetate. The organic layer was

washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 5% ethyl acetate in hexane to give title compound (1.328 g, 57%).

5


B.



A solution of Part A compound (200 mg, 0.55 mmol) and triphosgene (248 mg, 0.82 mmol) in pyridine (600  $\mu\text{L}$ ) was heated to  $160^\circ\text{C}$  in a sealed tube for 1 min. The reaction was cooled to room temperature and partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 25% ethyl acetate in hexane to give title compound (203 mg, 94%).

15


C.



Following the same experimental procedure as in Example 28 Part D, coupling between Part B compound and 3-amino phenyl boronic acid afforded the title compound (yield 99%).


20

D.




Following the same experimental procedure as in  
Example 28 Part G, alkylation of Part C compound with  
5 ethyl bromoacetate afforded the title compound (yield  
34%).

E.




10 Following the same experimental procedure as in  
Example 28 Part H, hydrolysis of Part D compound afforded  
the title compound (yield 90%).

### Example 30




15

A.



Following the same experimental procedure as in Example 28 Part D, the Suzuki coupling between Example 29 Part B compound and 3-methoxy phenyl boronic acid afforded the title compound (yield 87%).


B.



10


Following the same experimental procedure as in Example 28 Part E, Part A compound was treated with boron tribromide to afford the title compound (yield 73%).

15 C.




Following the same experimental procedure as in Example 28 Part G, alkylation of Part B compound with ethyl bromoacetate afforded the title compound (yield 96%).


D.



Following the same experimental procedure as in Example 28 Part H, hydrolysis of Part C compound afforded the title compound as a white solid (yield 98%).

Example 31

A.




10

Example 1 Part A compound was coupled with 2-formyl-phenyl boronic acid by Suzuki coupling procedure described in Example 1 Part B to give the title compound in 84% yeild.


15

B.




Triethyl phosphonoacetate (0.27 mL, 1.37 mmol) was added dropwise to a stirred suspension of sodium hydride (33 mg, 1.37 mmol) in THF (2 mL). The mixture was stirred for 1 hr, followed by dropwise addition of Part B compound (500 mg, 1.24 mmol) in THF (2 mL). The resulting yellow solution was stirred at room temperature overnight. Additional sodium hydride (20 mg, 0.83 mmol) was added and the reaction was stirred for 1 h. The reaction was diluted with water (50 mL) and ether (30 mL). The aqueous layer was extracted with ether twice. The combined organic layers were washed with brine dried over MgSO<sub>4</sub> and concentrated to give the title compound (528 mg, 90%) which was used without purification in the next step.


C.



A mixture of Part B compound (100 mg, 0.212 mmol) in THF (1 mL) was treated with 1N NaOH solution (0.424 mL, 0.424 mmol). After refluxing overnight, the mixture was concentrated, and diluted with ethyl acetate/water. The reaction mixture was acidified with 1N HCl to pH of 1. The resulting light yellow precipitate was filtered, washed with hexane and water. The solid was purified by flash chromatography on silica gel eluting with 1:1 hexane:ethyl acetate to give title compound (42 mg, 45%).


Example 32

A.




- 5           A suspension of Example 31 Part B compound (100 mg, 0.212 mmol) and 20% palladium on carbon (20 mg) in methanol was stirred under hydrogen (1 atm) overnight. The reaction mixture was filtered through a pad of Celite( 545 and the filtrate was concentrated. The
- 10   residue was purified by flash chromatography on silica gel eluting with a step gradient of 5% to 10% EtOAc in hexane to give title compound (92 mg, 92%).

B.




- 15           A mixture of Part A compound (90 mg, 0.190 mmol) in THF (0.5 mL) was treated with 1N NaOH solution (0.380 mL, 0.380 mmol). After stirring overnight at RT, the mixture was concentrated, diluted with water. The reaction
- 20   mixture was acidified with 1N HCl to pH of 1. The tan precipitate was filtered, washed with hexane and water. The solid was dried to give the title compound (60 mg, 71%).




Example 33

To a solution of Example 31 Part A compound (200 mg, 0.49 mmol) in THF (1.5 mL) and water (0.5 mL) at 0°C was added sulfamic acid (48 mg, 0.49 mmol) and sodium chlorite (45 mg, 0.49 mmol). The reaction was stirred for 1.5 h and diluted with ether. The organic layer was washed with water, brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude material was purified by preparative reverse phase HPLC to give the title compound (65 mg, 31%).

Example 34

15


A.



A mixture of 3-bromophenol (20 g, 116 mmol), K<sub>2</sub>CO<sub>3</sub> (31.95 g, 231.2 mmol) and DMF (120 mL) was treated with tert-butyl bromoacetate (20 mL, 231 mmol) and the mixture stirred for 16 h. The reaction was concentrated and the residue was dissolved in ethyl acetate. The organic


fraction was washed with water, brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by flash column chromatography on silica gel eluting with 5% ethyl acetate in hexane to give the title compound as a  
5 colorless oil (30 g, 100%).

B.




Part A compound was coupled with 2-formyl phenyl  
10 boronic acid by Suzuki coupling procedure described in  
Example 5 Part C to give the title compound in 69% yeild.


C.



15 A mixture of benzil (5.5 g, 26.4 mmol) and ammonium acetate (12.94 g, 168 mmol) in glacial acetic acid (120 mL) was treated with Part B compound (7.5 g, 24.0 mmol) and stirred at 120°C for 15 h. The reaction mixture was cooled, poured in water (300 mL) and precipitate was  
20 filtered. The solid was washed with hexane and dried to give the title compound as an off-white solid (12 g, 100%).


Example 35

A.



5           A solution of Example 34 compound (11.5 g, 22.8 mmol) in ethanol (300 mL) and sulfuric acid (3 mL) was stirred at room temperature overnight. The reaction was concentrated, diluted with ethyl acetate and neutralized with aqueous sodium carbonate solution. The organic  
10 layer was washed with water, aqueous sodium carbonate, and brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude was triturated with hexane and ether to give the title compound as a light yellow solid (9.13 g, 84%).

15           B.




A mixture of Part A compound (0.100 g; 0.210 mmol), K<sub>2</sub>CO<sub>3</sub> (0.058 g; 0.421 mmol) and DMF (1 mL) was treated

with allyl bromide (20  $\mu$ L, 0.231 mmol) and the mixture stirred for 16 h. Additional allyl bromide (20  $\mu$ L, 0.023 mmol) was added. The mixture was stirred for 2 h. The mixture was diluted with equal amounts of ethyl acetate and water. The organic fraction was washed with water, brine, dried ( $\text{MgSO}_4$ ) and concentrated. The residue was purified by flash column chromatography with a step gradient of 10% to 20% ethyl acetate in hexane to give the title compound as a colorless oil. (0.095 g, 88%).


10

C.




A mixture of Part B compound (0.095 g, 0.184 mmol) in methanol (0.5 mL) and THF (0.5 mL) was treated with 1N NaOH solution (0.276 mL, 0.276 mmol). After stirring overnight at RT, the mixture was concentrated, diluted with ethyl acetate. The reaction mixture was acidified with 1N HCl to pH of 1. The layers were equilibrated and separated. The organic fraction was washed with brine, dried over  $\text{MgSO}_4$ , and concentrated. The crude was recrystallized from ethyl acetate/hexane/dichloromethane to give the title compound as a white solid (0.032 g, 36%).

20


Example 36

A.



- 5           A mixture Example 5 Part A compound (3.75 g; 10.0 mmol) and sodium hydride (0.24 g, 10.0 mmol) in 35 mL of THF was stirred at RT for 15 min. The anion was slowly treated with methyl iodide (1.42 g, 10.0 mmol) and stirred overnight. The mixture was diluted with equal
- 10 amounts of water and ethyl acetate. The layers were equilibrated. The organic fraction was washed with water (2X20 mL), brine, dried over MgSO<sub>4</sub> and concentrated under vacuum to give a white solid. The solid was
- 15 The mother liquor was concentrated and recrystallized from methanol and a trace of water to give 1 g of additional material. The white solid was combined to give the title compound (3 g, 77%).


20           B.



A mixture of 3-bromophenol (5 g, 29.0 mmol), potassium carbonate (4.0 g, 29.0 mmol) and benzyl bromide (4.87 g, 28.5 mmol) in 40 mL of DMF was stirred at RT for


18 hr. The mixture was diluted with water, the pH adjusted to 12 with KOH (pellets) and a white solid formed. The solid was triturated with water, filtered and washed with water to give the title compound (7.20 g, 96%).

C.




A mixture of Part B compound (3.20 g, 12.16 mmol) in 24 mL of THF at -78°C was treated with n-butyllithium (5.8 mL, 14.6 mmol). After 1 h at -78°C the anion was slowly treated with triisopropyl borate (2.28 g, 12.18 mmol) and the mixture warmed to RT. The mixture was diluted with a 3% acetic acid/water solution (60 mL). The mixture was stirred for 40 min. and the solid which formed was collected. The solid was triturated with a small volume of 5% ethyl acetate/hexane to give the title compound as a white powder (0.5 g, 18%).

D.




Following the general experimental procedure as in Example 8 Part A, Suzuki coupling between Part A and Part C compound afforded the title compound (yield 60%).

E.




A mixture of Part D compound (0.50 g, 1.0 mmol) in ethanol (7 mL) was degassed and treated with 100 mg of 10% Pd on carbon. The mixture was put under an atmosphere of hydrogen gas (balloon pressure) and stirred overnight. The mixture was filtered and the colorless solution was mixed with 100 mg of Pearlman's catalyst and placed under an atmosphere of hydrogen for 4 h. The reaction was filtered and the filtrate concentrated to an oil. The oil was used without further purification.

F.




A mixture of the crude Part E compound (1 mmol), K<sub>2</sub>CO<sub>3</sub> (0.138 g, 1.0 mmol) and ethyl bromoacetate (0.14 g, 0.8 mmol) in dimethyl formamide (3 mL) was stirred at RT overnight. The contents were diluted with equal amounts of water and ethyl acetate. The organic fraction was dried (MgSO<sub>4</sub>) and concentrated to provide a thick oil. The oil was purified on silica gel by flash column chromatography with 15% ethyl acetate:hexanes to give 0.28 g of crude material. The material was rechromatographed on silica with 7% ethyl acetate/dichloromethane to give the title compound (0.18 g, 36%).


G.



The Part F compound (0.18 g, 0.36 mmol) was dissolved in ethanol (2 mL) and treated with 1N NaOH (2 mL, 2 mmol). After 18 h at RT the mixture was acidified with citric acid until a pH of 3 was achieved. The mixture was diluted with equal volumes of ethyl acetate and water. The layers were equilibrated and the organic fraction dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give the title compound as a (0.17 g, 100%).

Example 37

15 A.




A mixture of Example 5 Part A compound (0.75 g, 2 mmol) in 5 mL of DMF at RT was treated with K<sub>2</sub>CO<sub>3</sub> (0.28 g, 2 mmol) and n-propyl iodide (0.3 mL, 3 mmol). The mixture was stirred for 18 h and poured into water. The



water fraction was decanted from the thick residue. The residue was triturated with EtOH/water and the remainder was dried under vacuum to give the title compound as a white foam (380 mg, 50%).

5


B.



Nitrogen was bubbled through a solution of Part A compound (350 mg, 0.84 mmol), 3-amino phenyl boronic acid  
10 (172 mg, 1.26 mmol) and aqueous sodium carbonate (840  $\mu$ L, 2M, 1.68 mmol) in toluene (1.3 mL) and ethanol (700  $\mu$ L) at room temperature for 15 min.


Tetrakis(triphenylphosphine) palladium(0) (50 mg, 0.04 mmol) was added and the mixture was stirred at 80°C for  
15 36 h under argon. The reaction mixture was cooled to room temperature and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product  
20 was purified by flash chromatography on silica gel, eluting with 35:65 ethyl acetate/hexane to give title compound (310 mg, 88%).

C.



To a slurry of Part B compound (680 mg, 1.57 mmol) and potassium carbonate (420 mg, 3 mmol) in anhydrous dimethylformamide (3 mL) was added ethyl bromoacetate (250 mg, 1.50 mmol). The reaction mixture was stirred at room temperature for 18 h. The reaction was partitioned between diethyl ether and aqueous sodium bicarbonate solution. The organic layer was washed with water, and brine, then dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography eluting with 3:7 ethyl acetate/hexane to give title compound (450 mg, 55%).


15 D.



A mixture of Part C compound (0.35 g, 0.67 mmol) in methanol (3 mL) was treated with 1M NaOH solution (2 mL, 2 mmol). After stirring for 3 h at RT the mixture was diluted with ether and citric acid was added until the aqueous fraction maintained a pH of 3. The layers were equilibrated and separated. The organic fraction was


dried over  $\text{MgSO}_4$ , and concentrated to give the title compound (0.33 g, 100%).

Example 38



5


A.



Followed the same experimental procedure as described in Example 37 Part B using Example 5 Part A compound and 3-amino phenyl boronic acid to give the title compound (yield 50%).

10


B.




The experimental procedure as set out in Example 37 Part C employing the above Part B compound was followed to give the title compound (yield 55%).

15


C.



The experimental procedure as set out in Example 37  
Part D employing the above Part B compound was followed  
5 to give the title compound (yield 75%).

Example 39

A.




10

The experimental procedure as set out in Example 37  
Part B was followed for Suzuki coupling between the  
Example 36 Part A compound and 3-amino phenyl boronic  
acid to give the title compound (yield 76%).


15

B.



The experimental procedure as set out in Example 37 Part C employing the above Part B compound was followed to give the title compound (yield 75%).


5 C.




The experimental procedure as set out in Example 37 Part D employing the above Part B compound was followed to give the title compound (yield 60%).

10

### Example 40




A.



15 A mixture of Example 34 Part B compound (1.20 g, 3.84 mmol) in 10 mL of a 3:1 THF:water solution in ice bath was treated sequentially with sulfamic acid (1.31 g, 13.6 mmol) and sodium chlorite (1.22 g, 13.6 mmol). After 15 min the mixture was warmed to RT for 1 h and  
20 diluted with equal amounts of water and ether. The layers were equilibrated and the organic fraction washed

with brine, dried (MgSO<sub>4</sub>) and concentrated to give the title compound (1.20 g, 95%).

B.




5

A mixture of Part A compound (0.53 g, 1.61 mmol), potassium carbonate (0.23 g, 1.70 mmol) and ( $\alpha$ -bromoacetophenone (0.29 g, 1.70 mmol) in 4 mL of DMF was stirred at RT overnight. The mixture was diluted with equal amounts of water and ether. The layers were equilibrated. The organic fraction was washed with water (2 X 20 mL), brine, dried over MgSO<sub>4</sub> and concentrated under vacuum. The crude product was purified by flash chromatography on silica gel, eluting with 15:85 ethyl acetate/hexane to give title compound (590 mg, 82%).

15

C.




A mixture of Part B compound (0.59 g, 1.33 mmol) in 4 mL of acetic acid with ammonium chloride (0.60 g, 7.8 mmol) was heated to reflux for 48 h. The mixture was diluted with equal amounts of water and ethyl acetate. The organic fraction was washed with water, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by flash chromatography on silica gel eluting with 3:97:0.1

25


THF:dichloromethane:acetic acid to give title compound (70 mg, 14%).

Example 41



5


A.



A solution of Example 40 Part A compound (0.32 g, 1 mmol), 4-dimethylamino pyridine (0.04 g, 0.3 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.23 g, 1.20 mmol) and 2-pyridoin (0.21 g, 1.0 mmol) in dichloromethane (3 mL) was stirred at room temperature for 4 hr. The reaction mixture was washed with water, brine, dried over anhydrous  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash chromatography on silica gel, eluting with 3:7 ethyl acetate/hexane to give title compound (0.096 g, 20%).

15

B.




20


A mixture of Part A compound (0.096 g, 0.18 mmol) in 2 mL of acetic acid with ammonium chloride (0.096 g, 1.2

mmol) was heated to reflux for 4 h. The mixture was stripped under reduced pressure to an oil. The oil was purified on reverse phase column chromatography. The pure fractions were combined and concentrated to give the  
5 title compound (10 mg, 13%).

Example 42



A.



10


A solution of 2-iodo-4-chlorobenzoic acid (3.5 g, 12.4 mmol), 4-dimethylaminopyridine (1.65 g, 13.5 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.6 g, 13.5 mmol) and benzoin (2.63 g, 12.4 mmol) in dichloromethane (20 mL), was stirred at room temperature for 4 h. The reaction mixture was washed with HCl (1N), NaOH (0.1 N), brine, dried over anhydrous MgSO<sub>4</sub> and concentrated to give the crude ketoester (5.9 gm, 99%).

20

A mixture of ketoester (5.9 g, 12 mmol) and ammonium chloride (5.50 g, 71 mmol) in glacial acetic acid (45 mL) was stirred at reflux for 4 hr. The reaction mixture was cooled and poured on ice. The tar-like solid was digested with hot ethanol (25 mL). The resulting beige  
25 solid was collected to give the title compound (4.5 g, 88%).




B.




The experimental procedure as described in Example  
5 37 Part B was followed employing the above Part A  
compound to give the title compound (yield 92%).

C.



10 The experimental procedure as described in Example  
37 Part C was followed, employing the above Part B  
compound, to give the title compound (yield 80%).

D.




15


The Part C compound (0.48 g, 0.94 mmol) was diluted  
with ethanol (8 mL) and treated with NaOH (0.40 g; 10  
mmol) and 3 mL of water. After 18 h at RT the mixture  
was acidified with citric acid to pH 3. The mixture was

diluted with equal volumes of ethyl acetate and water. The layers were equilibrated and the organic fraction dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The residue was purified by preparative reverse phase column chromatography. The isolated material was further purified by flash column chromatography on silica gel with gradient elution from 4% ethanol in dichloromethane to 10% ethanol in dichloromethane with a trace of acetic acid to give the title compound (90 mg, 20%).

10

Example 43


A.



15 A mixture of cyclohexane carboxaldehyde (2.800 gm, 25 mmol) and trimethylsilyl cyanide (2.475 gm, 25 mmol) was treated with zinc iodide (2 mg, 0.006 mmol). The mixture was stirred at RT for 3 h. The title compound was used without further purification, yield 100%.

20

B.




A mixture of Part A compound (2.00 g, 9.45 mmol) in THF (20 mL) at RT was treated with phenylmagnesium chloride (3M in THF, 6.6 mL, 20 mmol). The mixture was stirred for 1 h at RT and then heated to reflux for 4 h.

5 The reaction mixture was cooled and stirred with an excess of 1N HCl solution for 18 h. The mixture was partitioned between ether and water. The organics were dried over MgSO<sub>4</sub>, concentrated, and the residue purified by flash column chromatography on silica gel eluting with

10 a step gradient of 1:9 to 3:7 ethyl acetate/hexane to give the title compound (750 mg, 34%).

C.




15 A solution of Example 40 Part A compound (330 mg, 1.0 mmol), 4-dimethylamino pyridine (40 mg, 0.3 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (230 mg, 1.2 mmol) and Part B compound (220 mg, 1.0 mmol) in dichloromethane (2 mL) was stirred at room temperature

20 for 4 h. The reaction mixture was washed with water (2x50 mL), brine (50 mL), dried over anhydrous MgSO<sub>4</sub> and concentrated. The residue was purified by flash column chromatography on silica gel eluting with a step gradient of 5:95 to 10:90 ethyl acetate/hexane to give title


25 compound (400 mg, 75%).

D.




A mixture of Part C compound (400 mg, 0.75 mmol) and ammonium acetate (580 mg, 3.1 mmol) in glacial acetic acid (4 mL) was stirred at reflux for 18 h. The reaction mixture was cooled and concentrated. The crude oil was purified by flash column chromatography on silica gel eluting with 3:97:0.5 methanol/dichloromethane/acetic acid to give the title compound (270 mg, 78%).

10

Example 44

A.




A mixture of Example 43 Part A compound (2.11 g, 10 mmol) in THF (10 mL) at RT was treated with cyclohexylmagnesium chloride (2M in ether, 8 mL, 16 mmol). The mixture was stirred for 1 h at RT and then heated to reflux for 3 h. The reaction mixture was cooled and stirred with an excess of 1N HCl solution for 4 h. The mixture was partitioned between ether and water. The organics were dried over MgSO<sub>4</sub>, concentrated,

20

and the residue purified by flash column chromatography on silica gel eluting with a step gradient of 1:9 to 4:6 ethyl acetate/hexane to give title compound as a foam (750 mg, 33%).


5

B.




Part A compound was coupled with Example 40 Part A compound employing the procedure described in Example 43  
10 Part C to obtain the title compound in 18% yield.

C.




A mixture of Part B compound (240 mg, 0.45 mmol)  
15 and ammonium chloride (240 mg, 3.1 mmol) in glacial acetic acid (4 mL) was stirred at reflux for 24 h. The reaction mixture was cooled poured in water and filtered. The crude beige solid was triturated with water to get the title compound (180 mg, 87%).

20

Example 45

A.




5        A mixture of Example 5 Part A compound (0.75 g, 2 mmol) in 5 mL of DMF at RT was treated with  $K_2CO_3$  (0.28 g, 2 mmol) and bromoethyl acetate (1 mL, 6 mmol). The mixture was heated to 100°C in a sealed tube stirred for 72 h. The mixture was cooled to RT and poured into

10    water. The organics were extracted with ethyl acetate, dried over  $Na_2SO_4$  and concentrated. The residue was purified on flash column chromatography with 15:85 ethyl acetate/hexane to give the title compound (0.40 g, 46%).

15


B.



Nitrogen was bubbled through a solution of Part A compound (300 mg, 0.69 mmol), 3-aminophenyl boronic acid

(158 mg, 1.00 mmol) and aqueous sodium carbonate (1 mL, 2M, 2 mmol) in toluene (2 mL) and ethanol (1 mL) at room temperature for 15 min. Tetrakis(triphenylphosphine) palladium(0) (550 mg, 0.04 mmol) was added and the  
5 mixture was stirred at 80°C for 36 hr under argon. The reaction mixture was cooled to room temperature and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified  
10 by flash chromatography on silica gel, eluting with 4:6 ethyl acetate/hexane to give title compound (150 mg, 46%).


C.




15

To a slurry of Part B compound (120 mg, 0.28 mmol) and potassium carbonate (40 mg, 0.28 mmol) in anhydrous dimethylformamide (3 mL) was added ethyl bromoacetate (50 mg, 0.28 mmol). The reaction mixture was stirred at room  
20 temperature for 18 h. The reaction was partitioned between diethyl ether and aqueous sodium bicarbonate solution. The organic layer was washed with water, and brine, then dried over anhydrous MgSO<sub>4</sub> and concentrated. The crude product was purified by flash chromatography  
25 eluting with 3:7 ethyl acetate/hexane to give title compound (95 mg, 60%).

D.




A mixture of Part C compound (90 mg, 0.16 mmol) in methanol (3 mL) was treated with 1M NaOH solution (1 mL, 1 mmol). After stirring for 3 h at reflux the mixture was cooled and diluted with ether. Citric acid was added until the aqueous fraction maintained a pH of 3. The layers were equilibrated and separated. The organic fraction was dried over  $\text{MgSO}_4$ , and concentrated to give the title compound (80 mg, 100%).

Example 46


A mixture of Example 40 Part B compound (0.59 g, 1.33 mmol) in 4 mL of acetic acid with ammonium chloride (0.60 g, 7.8 mmol) was heated to reflux for 48 h. The mixture was diluted with equal amounts of water and ethyl acetate. The organic fraction was washed with water, dried ( $\text{MgSO}_4$ ) and concentrated. The remainder was purified on silica by flash column chromatography with 3:97:0.1 THF:dichloromethane:acetic acid to give 15 mg (3%) title compound.



### Example 47




A.




5 A mixture of Example 5 Part A compound (0.75 g, 2  
mmol) in DMF (5 mL) was treated with K<sub>2</sub>CO<sub>3</sub> (420 mg, 4  
mmol) and isobutyl iodide (740 mg, 4 mmol). The mixture  
was heated in a sealed tube at 60°C for 18 h. The tube  
was cooled, and the contents diluted with equal amounts  
10 of water and ether. The organic fraction was dried over  
anhydrous MgSO<sub>4</sub> and concentrated to a thick oil. The  
crude product was purified by flash chromatography on  
silica gel, eluting with a step gradient of 2:98 to 7:93  
ethyl acetate/hexane to give title compound (250 mg,  
15 30%).

B.



The experimental procedure as set out in Example 37 Part B employing the above Part A compound was followed to give the title compound (yield 62%).


5 C.



The experimental procedure as set out in Example 37 Part C employing the above Part B compound was followed to give the title compound (yield 72%).


10

D.




The experimental procedure as set out in Example 37 Part D employing the above Part C compound was followed to give the title compound (yield 61%).

### Example 48




A.



A solution of Example 6 Part A compound (0.49 g, 1.18 mmol) in DMF (2 mL) at RT was treated with  $K_2CO_3$  (0.28 g, 2 mmol) and bromoacetonitrile (0.14 g; 1.20 mmol). The mixture was stirred overnight at RT and then heated to 60°C for 4 h. The reaction mixture was cooled to RT and poured into water. The solid was collected and dried overnight to give the title compound (0.45 g, 85%).

10


B.



A solution of Part A compound (0.40 g, 0.88 mmol) in xylene (2 mL) was treated with azidotrimethyltin (0.23 g, 1.14 mmol) and the mixture was heated under  $N_2$  for 18 h at 135°C. The reaction was quenched with methanol and the mixture concentrated to an oil. The oil was purified by flash column chromatography on silica with a step wise gradient using 15/85 ethyl acetate/hexane then 5/95 methanol/dichloromethane to give 0.30 g of the title compound as a white foam (68%).

15


20

Example 49

A mixture of Example 34 Part B compound (0.20 g, 0.64 mmol), pyridinium *p*-toluenesulfonate (40 mg), and  
5 meso-1,2-dihydroxy-1,2-diphenylethane (0.14 g, 0.64 mmol) in toluene (5 mL) was heated to 80°C for 36 h and then brought to reflux for 3 h. The mixture was cooled to RT and diluted with water and ethyl acetate. The organics were dried (MgSO<sub>4</sub>) and carried on to the next step  
10 without further characterization.

The crude ester (~ 0.64 mmol) in methanol (4 mL) was treated with 1M NaOH (1 mL, 1 mmol) and stirred overnight. The mixture was concentrated and the residue purified by chromatography on SP207 gel with a stepwise  
15 gradient from water to 80% methanol/water in 20% increments (70 mL portions). The pure fractions were combined and freeze dried to give the title compound as a white lyophilate (130 mg, 45%).

20


Example 50

A mixture of Example 34 Part B compound (0.20 g, 0.64 mmol), pyridinium *p*-toluenesulfonate (40 mg), and  
(*R,R*)-1,2-dihydroxy-1,2-diphenylethane (0.14 g, 0.64  
25 mmol) in toluene (5 mL) was heated to 80°C for 36 h and then brought to reflux for 3 h. The mixture was cooled

to RT and diluted with water and ethyl acetate. The organics were dried ( $\text{MgSO}_4$ ) and carried on to the next step without further characterization.


The crude ester (~ 0.64 mmol) in methanol (4 mL) was treated with 1M NaOH (1 mL, 1 mmol) and stirred overnight. The mixture was concentrated and the residue purified by chromatography on SP207 gel with a stepwise gradient from water to 80% methanol/water in 20% increments (70 mL portions). The pure fractions were combined and freeze dried to give the title compound as a white lyophilate (200 mg, 65%).

#### Example 51



15


A.



20


The experimental procedure as set out in Example 37 Part A using *tert*-butyl bromoacetate as the alkylating agent was followed to give the title compound (yield 92%).

B.




The experimental procedure as set out in Example 37 Part B employing the above Part A compound was followed to give the title compound (yield 76%).

C.



The experimental procedure as set out in Example 37 Part C employing the above Part B compound was followed using *tert*-butyl bromoacetate to give the title compound (yield 75%).

D.




15


Part C compound (0.16 g, 0.26 mmol) in a 1:1 mixture of trifluoroacetic acid/ dichloromethane (4 mL) was stirred at room temperature overnight. The mixture was concentrated and the residue adjusted to pH 13 with 1N NaOH solution. The solution was purified by chromatography on SP207 gel with a stepwise gradient from

water to 60% methanol/water in 20% increments (80 mL portions). The pure fractions were combined, the pH of the combined fractions were adjusted to pH 6.5 with 0.1 N NaOH and freeze dried to give 83 mg (62%) of title  
5 compound as a white lyophilate.

Example 52




A.




10

A mixture of Example 8 Part B compound (0.25 g, 0.6 mmol), K<sub>2</sub>CO<sub>3</sub> (0.12 g, 0.9 mmol) and DMF (3 mL) was treated with bromoacetonitrile (0.11 g; 0.9 mmol) and the mixture stirred for 72 h. TLC indicated that the  
15 reaction was not complete and additional bromoacetonitrile (0.11 g, 0.9 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.12 g, 0.9 mmol) were added. The mixture was stirred for 18 h. The mixture was diluted with equal amounts of ethyl acetate and water. The organic fraction was washed with  
20 water, dried (MgSO<sub>4</sub>) and concentrated. The remainder was purified by flash column chromatography with 20% ethyl acetate in hexane to give 0.22 g (80%) of the title compound.

B.




A mixture of Part A compound (0.20 g, 0.44 mmol) in xylenes (2 mL) was treated with azidotrimethyltin (0.11 g, 0.53 mmol) and warmed to 135°C (bath temp) for 18 h. The mixture was concentrated and stirred with methanol for 45 min. The methanol solution was concentrated and the residue purified by flash column chromatography with a step gradient elution using dichloromethane (80 mL), 1% methanol/dichloromethane (200 mL) and 2% methanol/dichloromethane (200 mL) to give 36 mg (16%) of the title compound.

Example 53

15

A.



A solution of 1,3-diphenyl-1,3-propanedione (4.50 g, 20 mmol) in 150 mL of dichloromethane was mixed with triethylamine (2.07 g, 20 mmol) and cooled to -5°C (internal temperature). The mixture was then treated with 4-aetamidobenzenesulfonyl azide (4.80 g, 21 mmol). The reaction was maintained at -4°C for 4 h and then




warmed to RT overnight. The mixture was diluted with water and the organic fraction was washed with 1 N NaOH solution (4 X 125 mL), water (2 X 50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to an oil. The oil was diluted with a small volume (15 mL) of methanol and 3 drops of water. The resulting solid was collected and dried to give 1.50 g (30%) of the title compound.

(Reference Meier, H. Chem Ber. 119, 3382-3393, 1986.)

10


B.



A mixture of the Part A compound (1.5 g; 6 mmol) in 50 mL of ethanol was refluxed for 18 h. The ethanol was stripped off under reduced pressure to leave a thick oil. The oil was solidified by mixing with hexane (10 mL) and concentrating under reduced pressure. After drying 1.6 g (100%) of the title compound was obtained as a beige solid.

20

C.




A mixture of Part B compound (0.1 g, 0.37 mmol) in 3 mL of ethanol was treated with 2-bromophenylhydrazine hydrochloride (84 mg, 0.37 mmol) and heated to reflux overnight. The ethanol was removed and the remainder was purified on silica gel by flash column chromatography

25

with 65:35 ethyl acetate:hexane to give 90 mg (62%) of the title compound.


D.



5 A mixture of Part C compound (0.13 g, 0.33 mmol),  $K_2CO_3$  (0.10 g, 0.73 mmol) and DMF (3 mL) was treated with methyl iodide (0.10 g, 0.70 mmol) and the mixture stirred for 18 h. The mixture was diluted with equal amounts of  
10 ethyl acetate and water. The organic fraction was washed with water, dried ( $MgSO_4$ ) and concentrated. The remainder was purified by flash column chromatography with 10% ethyl acetate in hexane to give 80 mg (60%) of the title compound.

15


E.



A mixture of Part D compound (80 mg, 0.19 mmol), toluene (1 mL), ethanol (0.5 mL), 3-aminophenyl boronic acid (35 mg, 0.25 mmol), and 2 N  $Na_2CO_3$  solution (0.125  $\mu$ L, 0.25 mmol) was degassed with a gentle  $N_2$  flow for 10  
20 min. The mixture was treated with tetrakis(triphenylphosphine)-palladium (0) (10 mg) and the mixture heated to 80°C for 18 h. The mixture was cooled  
25 to RT and diluted with equal portions of water and ethyl acetate. The organic fraction was dried over  $MgSO_4$  and concentrated. The remainder was purified on silica gel by flash column chromatography with 65:35 ethyl

acetate:hexane as a mobile phase to give 66 mg (80%) of the title compound.

F.




5

A mixture of DMF (0.5 mL), potassium carbonate (30 mg, 0.21 mmol) and Part E compound (60 mg, 0.15 mmol) was treated with ethyl bromoacetate (17  $\mu$ L, 0.15 mmol). The mixture was stirred overnight and diluted with equal portions of ethyl acetate and water. The layers were equilibrated, the organic fraction dried ( $\text{MgSO}_4$ ) and concentrated. The residue was purified on silica gel by flash column chromatography with 20:80 ethyl acetate:hexane as a mobile phase to give 60 mg (80%) of the title compound.

15


G.




20

A solution of Part F compound (60 mg, 0.12 mmol) and methanol (3 mL) was treated with 1 N NaOH solution (1 mL, 1 mmol) and stirred overnight. The mixture was acidified with citric acid until a pH of 3 was maintained. The resulting slurry was diluted with equal portions of ethyl acetate and water and the layers equilibrated. The organic fraction was dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Pumping overnight gave 50 mg (88%) of the title compound.


25

Example 54

The title compound was generated by the procedure  
5 described in Example 19 starting with Compound A Example  
13.


Example 55

10 The title compound was generated following the  
procedure described in Example 34 utilizing 4,4'-  
difluorobenzil.


Example 56

15


The title compound was generated from Example  
following the procedure described in Example 35.

Example 57


Following the procedure described in Example 35,  
Compound A Example 35 was alkylated with chloromethyl  
5 methyl ether and hydrolyzed to give the title compound.

Example 58


Following the procedure described in Example 35,  
10 Compound A Example 35 was alkylated with 1-bromo-2-  
fluoroethane and hydrolyzed to give the title compound.

Example 59


15 The title compound was generated by the procedure  
described in Example 8 starting with Example 19 Part E  
compound.

Example 60

The title compound was generated following the procedure described in Example 23, starting with Example 21 Part A compound.

Example 61

A.




10

To a stirred solution of 2-bromobenzoyl chloride (1.31 mL, 10.0 mmol) in THF (10 mL) under argon at -22°C was added tributylphosphine (2.74 mL, 11.0 mmol) over 10 min. The temperature was not allowed to rise above -15°C. After 20 min, a solution of n-propylmagnesium chloride (5.0 mL, 10.0 mmol, 2 M in ether) was added in one portion. The reaction temperature rose almost at once to 34°C and subsided to -20°C over 10 min. After an additional 10 min, 1 M aqueous HCl (18 mL) was added and the mixture was extracted with ether (100 mL). The ether extract was washed once with brine, once with saturated sodium bicarbonate solution, dried (MgSO<sub>4</sub>) and evaporated. Purification by flash chromatography on

silica gel (5x25 cm column, 38:62 CH<sub>2</sub>Cl<sub>2</sub>/hexanes) gave the title compound as a yellow oil, 2.28 g, 100% yield.

B.




5

To a stirred solution of potassium hydroxide (1.23 g, 22 mmol) in methanol (10 mL) at room temperature under argon was added Part A compound (1.00 g, 4.40 mmol). After five min, benzaldehyde (1.34 mL, 13.2 mmol) was added in one portion. After 4 h, the reaction mixture was cooled to 0°C, treated with 1 M aqueous HCl to bring the solution to pH 7.5 and then evaporated at < 30°C. The residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and brine. The aqueous phase was extracted twice with CH<sub>2</sub>Cl<sub>2</sub>, and the extracts were combined, dried (MgSO<sub>4</sub>) and evaporated. Purification by flash chromatography on silica gel (5x15 cm column, 2:3 CH<sub>2</sub>Cl<sub>2</sub>/hexanes) gave the title compound as a light yellow oil, 1.29 g, 95% yield.

20


C.



To a stirred slurry of benzaldehyde (5.1 mL, 50.0 mmol), potassium cyanide (13 g, 200 mmol), and t-butyldimethylsilyl chloride (9.0 g, 60 mmol) in acetonitrile (100 mL) at room temperature under argon was added zinc iodide (250 mg, 0.8 mmol). After 14 h, the reaction mixture was filtered and the filtrate evaporated. The evaporate was slurried in hexanes for 1 h under nitrogen, refiltered and evaporated. The oily residue was distilled (80-83°C, 0.8 torr) to give the title compound as a colorless oil, 7.68 g, 62% yield.


30

D.



5           A solution of lithium diisopropylamide was prepared at 0°C under argon from diisopropylamine (1.2 mL, 8.1 mmol) and n-butyllithium (3.2 mL, 8.0 mmol, 2.5 M in hexanes) in ether (20 mL). To this solution at -72°C was added a solution of Part C compound (1.80 g, 7.30 mmol) in ether (5 mL). The temperature was not allowed to rise above -67°C. After 1 h, a solution of Part B compound (2.30 g, 7.30 mmol) was added over the course of 10 min. The reaction was stirred at -78°C for 2 h and then stored at -40°C for 14 h. The reaction was then warmed to 0°C for 30 min and then quenched with saturated ammonium chloride solution. The reaction mixture was extracted with ether (100 mL). The extract was dried (MgSO<sub>4</sub>) and evaporated. Purification by flash chromatography on silica gel (5x25 cm column, 1:1 CH<sub>2</sub>Cl<sub>2</sub>/hexanes) gave the title compound as a white amorphous solid, 1.88 g, 46% yield.

E.




25

To a solution of Part D compound (1.65 g, 2.93 mmol) in THF (5 mL) at room temperature under argon was added tetrabutylammonium fluoride solution (3.6 mL, 3.6 mmol, 1 M in THF). After 1 h, the reaction was quenched with saturated sodium bicarbonate solution and extracted twice with CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The extracts were combined, dried (MgSO<sub>4</sub>) and evaporated. Purification by flash




chromatography on silica gel (5x25 cm column, 1:1 CH<sub>2</sub>Cl<sub>2</sub>/hexanes) gave the title compound (as a mixture of diastereomers) as a colorless oil, 1.05 g, 85% yield.

5 F.




Analogous to the procedure described in Example 19 Part C, Compound E (600 mg, 1.42 mmol) gave the title compound as a colorless oil, 557 mg, 97% yield.

G.




Analogous to the procedures described in Example 8, Part F compound gave the title compound.

#### Example 62




Analogous to the procedure described in Example 19, Example 21 Part A compound gave the title compound.

Example 63


Analogous to the procedure described in Example 8,  
Example 21 Part A compound gave the title compound.

5

Example 64


Analogous to the procedure described in Example 8,  
Example 20 Part A compound gave the title compound.

10


Example 65

Analogous to the procedure described in Example  
23, Example 61 Part F compound gave the title compound.

15


Example 66

A.



To a stirred slurry of NaH (60% oil dispersion, 10.0 g, 250 mmol) in diethyl carbonate (40 mL) and toluene (15 mL) at room temperature under argon, was added absolute ethanol (150  $\mu$ L) and then a solution of 2-bromoacetophenone (24.88 g, 125 mmol) in toluene (20 mL) over 20 min. The reaction temperature rose autogenously to 80°C and then was maintained at that temperature for 2h. The mixture was cooled to room temperature and acetic acid (15 mL) was added. The resulting bulky solid was treated with ice-cold water (100 mL) and then extracted twice with ether (250 mL). The extracts were combined, dried ( $\text{MgSO}_4$ ) and evaporated to give the crude ketoester as a yellow oil, 29.75 g. Purification of a portion of the crude product (10g) by flash chromatography on silica gel (5x25 cm column, 9:1  $\text{CH}_2\text{Cl}_2$ /hexanes) gave the title compound as a light yellow oil, 7.83 g, 70% calculated yield.

B.




A solution of Part A compound (5.42 g, 20.0 mmol), benzaldehyde (2.24 mL, 22 mmol), 6-aminocaproic acid (100 mg, 0.76 mmol) and acetic acid (3 mL) in benzene (10 mL) was heated to reflux for 1 h using a Dean-Stark trap. The reaction mixture was cooled to room temperature, diluted with ethyl acetate and washed four times with saturated sodium bicarbonate solution until the washings were pH 8. The organic phase was dried ( $\text{MgSO}_4$ ) and evaporated. Purification of the crude product by flash

chromatography on silica gel (5x25 cm column, 7:3 CH<sub>2</sub>Cl<sub>2</sub>/hexanes) gave the title compound (55:45 mixture of geometric isomers) as a white amorphous solid, 4.38 g, 61% yield.

5


C.



Analogous to the procedure described in Example 19 Part B, Part B compound was used to prepare the title compound.

10


D.



Analogous to the procedure described in Example 19 Part C, Part C compound was used to prepare the title compound.

15


E.



20


Analogous to the procedure described in Example 8 Part A, Part D compound was used to prepare the title compound.

F.




Analogous to the procedure described in Example 8  
5 Part B, Part E compound was used to prepare the title  
compound.

G.




10 To a stirred solution of Example 66 Part F  
compound (497 mg, 1.08 mmol) in THF (2 mL) at room  
temperature under argon was added a solution of lithium  
aluminum hydride (1.5 mL, 1.5 mmol, 1 M in THF) over 30  
sec. The resulting light yellow solution was stirred for  
15 15 min, whereupon a gelatinous mixture formed. This was  
quenched cautiously with 5% potassium hydrogen sulfate  
solution and extracted three times with ethyl acetate (20  
mL). The organic extracts were combined, dried (MgSO<sub>4</sub>)  
and evaporated to give a light pink solid, 440 mg, 97%  
20 yield.

H.




Analogous to the procedure described in Example 8  
 Part C, Part G compound was used to prepare the title  
 5 compound.


I.



By the procedure described in Example 8 Part D,  
 10 Part H compound was used to prepare the title compound.


Example 67

A.




To a stirred solution of Example 66 Part H compound (112 mg, 0.222 mmol) in methanol (5 mL) was added concentrated sulfuric acid (2 drops). The mixture was heated in a sealed tube at 100°C for 9 h. The reaction mixture was cooled to room temperature, treated with saturated sodium bicarbonate solution and evaporated to dryness. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, dried (MgSO<sub>4</sub>) and evaporated to give the title compound as a colorless oil, 98 mg, 88% yield.

B.




Analogous to the procedure described in Example 8 Part D, Part A compound was used to prepare the title compound.


Example 68



By the procedure described in Example 67, Example 66 Part H compound was used to prepare the title compound, substituting ethanol for methanol in the procedure.


Example 69

A.



5            Analogous to the procedure described in Example 20, Example 61 Part E compound was used to prepare the title compound.


B.



10


          Analogous to the procedure described in Example 8, Example 69 Part A compound was used to prepare the title compound.

15

Example 70




A.



A steady stream of dried air was bubbled into a stirred slurry consisting of Example 66 Part H compound (504 mg, 1.0 mmol), powdered 4 Å molecular sieves (200 mg) and tetrapropylammonium perruthenate (30 mg, 0.085 mmol) in toluene (3.5 mL) as the reaction was heated to 65°C. After 1 h, the reaction was cooled to room temperature, diluted with CH<sub>2</sub>Cl<sub>2</sub> and filtered through silica gel (~20 g). Evaporation gave the title compound as a colorless oil, 490 mg, 97%).


B.




15

To a solution of Part A compound (125 mg, 0.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> at room temperature under argon, was added a solution of trimethylaluminum (0.25 mmol, 0.5 mmol, 2 M in CH<sub>2</sub>Cl<sub>2</sub>), dropwise over 5 min. After 1 h, the reaction was quenched cautiously with 10% citric acid solution and extracted once with CH<sub>2</sub>Cl<sub>2</sub>. The organic phase was dried (MgSO<sub>4</sub>) and evaporated. Purification of the crude product by flash chromatography on silica gel (2.5x15 cm column, 3:97 ether/ CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound as a colorless oil, 124 mg, 96% yield.

C.




5      Analogous to the procedure described in Example 8  
Part D, Part B compound was hydrolyzed to prepare the  
title compound.

Example 71


10

Analogous to the procedure described in Example 23,  
Example 69 Part A compound was used to prepare the title  
compound.


15

Example 72

20      Analogous to the procedure described in Example 3,  
Example 61 Part F compound was used to prepare the title  
compound.

Example 73

A.




5

To a stirred slurry of 2-bromophenylhydrazine hydrochloride (31.02 g, 0.139 mol) at room temperature under nitrogen in 95% ethanol (500 mL) was added sodium acetate trihydrate (18.87 g, 0.139 mol) and deoxybenzoin (30.96 g, 0.139 mol). The mixture was heated to reflux for 2 h and then partially cooled and evaporated. The residue was slurried in saturated sodium bicarbonate solution (300 mL), filtered, washed with water and dried in vacuo at 60°C for 16 h. The resulting solids were triturated in hot methanol (200 mL), filtered and dried to provide the title compound (96:4 mixture of Z/E isomers) as a white solid, mp 126-128°C, 36.84 g, 76% yield.

20


B.



To a stirred slurry of sodium hydride (60% dispersion, 1.43 g, 35.8 mmol) in N-methylpyrrolidone (NMP, 25 mL) at room temperature under nitrogen, was added a solution of Part A compound (10.0 g, 27.4 mmol)

in NMP (15 mL) over 5 min. The temperature autogenously rose to 54°C as gas was evolved and a deep red solution formed. The reaction mixture was heated to 60°C for 3 h and then cooled to room temperature. To this solution  
5 was added freshly distilled propionic anhydride (4.6 mL, 35.9 mmol) at a rate to keep the reaction temperature below 35°C. The reaction mixture was stirred 2 h, then quenched with 5% potassium hydrogen sulfate solution and extracted four times with ether (100 mL). The ether  
10 extracts were combined, washed twice with water, once with brine, dried (MgSO<sub>4</sub>) and evaporated to give an orange oil, 12.9 g. Purification of the crude product by flash chromatography on silica gel (12x30 cm column, 1:4 ethyl acetate/hexanes) gave the title compound as a light  
15 yellow oil, 8.15 g, 71% yield.


C.



20 To a solution of Part B compound (2.33 g, 5.53 mmol) in DMF (15 mL) at room temperature under nitrogen was added sodium hydride (60% dispersion, 550 mg, 13.8 mmol) in several portions over 1 min. The reaction warmed autogenously to 38°C and was further heated to  
25 50°C for 20 min. The deep red reaction mixture was cooled in a water ice bath and quenched by adding 5% potassium hydrogen sulfate solution (10 mL) dropwise over 10 min. The mixture was then acidified with 1 N hydrochloric acid to bring the mixture to pH 2 and then  
30 extracted three times with ether (50 mL). The combined organic extracts were washed twice with water and once with brine, dried (MgSO<sub>4</sub>) and evaporated. Purification of the crude product by flash chromatography on silica gel (5x25 cm column, 1:3 ethyl acetate/hexanes) gave the

title compound as a light yellow amorphous solid, 1.63 g, 73% yield.

D.




5

The title compound was generated by the procedure described in Example 8 starting with Compound C Example 73.

10


Example 74



The title compound was generated by the procedure described in Example 23 starting with Example 73 Part C compound.


15

Example 75




20

The title compound was generated by the procedure described in Example 3 starting with Example 73 Part C compound.

Example 76

A.




5


A stirred slurry of sodium hydride (60% dispersion, 1.64 g, 41.1 mmol) in N-methylpyrrolidone (NMP, 13 mL) at room temperature under nitrogen, was heated to 40°C for 30 min. To this mixture was added Example 73 Part A compound (5.00 g, 13.7 mmol) in NMP (30 mL) in one portion. The reaction mixture was heated to 60°C for 4 h and then cooled to room temperature. To this solution was added freshly distilled acetyl chloride (4.3 mL, 57.5 mmol) in NMP (5 mL) at a rate to keep the reaction temperature below 50°C. The reaction mixture was stirred 30 min, then quenched with water (300 mL), extracted twice with ether (100 mL) and twice with ethyl acetate. The organic extracts were combined, washed with saturated sodium bicarbonate solution, water, and brine, dried (MgSO<sub>4</sub>) and evaporated. Purification of the crude product by flash chromatography on silica gel (5x20 cm column, 1:2 hexanes/CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound as a white amorphous solid, 1.20 g, 23% yield.

25


B.



The title compound was generated by the procedure  
5 described in Example 8 starting with Part A compound.


Example 77

10 The title compound was generated by the procedure  
described in Example 23 starting with Example 76 Part A  
compound.

Example 78

15

A.




20 A mixture of 2-bromoaniline (2.18 g, 12.7 mmol)  
and 2,3-diphenylmaleic anhydride (3.17 g, (12.7 mmol)  
were stirred and heated under argon to 200°C for 6 h.  
The mixture was cooled to room temperature, dissolved in

CH<sub>2</sub>Cl<sub>2</sub> (50 mL), dried (MgSO<sub>4</sub>) and evaporated.

Purification of the crude product by flash chromatography on silica gel (5x25 cm column, 2 L of 3:7 hexanes/CH<sub>2</sub>Cl<sub>2</sub>, then CH<sub>2</sub>Cl<sub>2</sub>) gave first 4-(2-bromoanilino)-2,3-diphenyl-4-

5 hydroxybutyrolactone (2.16 g, 40% yield) as a yellow solid, mp 162-164°C and then the title compound as a bright yellow solid, mp 178-180°C, 2.00 g, 39% yield.

B.




10

The title compound was generated by the procedure described in Example 8 starting with Example 78 Part A compound.


15

Example 79



20

A.




25

Analogous to the procedure described in Example 23 Part A, Example 73 Part C compound was used to prepare the title compound.




B.




5 To a stirred solution of triethylphosphonoacetate (0.50 mL, 2.5 mmol) in THF (2.5 mL) under nitrogen at room temperature was added sodium hydride (60% dispersion, 100 mg, 2.5 mmol). After 10 min, the reaction was heated to 50°C. After an additional 20 min, a solution of Example 73 Part C compound (775 mg, 1.80 mmol) in THF (2 mL) was added as one portion. The reaction mixture was stirred for 2 h, cooled to room temperature and partitioned between ethyl acetate (50 mL) and 5% potassium hydrogen sulfate solution. The organic phase was dried (MgSO<sub>4</sub>) and evaporated. Purification of the crude product by flash chromatography on silica gel (5x15 cm column, 1:49 ether/CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound as a colorless oil, 785 mg, 88% yield.


20 C.



Analogous to the procedure described in Example 8 Part D, Part B compound was used to prepare the title compound.

Example 80

A.




5

Hydrogenation of Example 79 Part B compound (765 mg, 1.53 mmol) was performed at room temperature and atmospheric pressure in ethanol (10 mL) using 10% palladium/charcoal (100 mg). After 4 h, the reaction mixture was purged with nitrogen, filtered (0.45  $\mu$  nylon filter) and the filtrate evaporated to provide the title compound as a colorless oil, 770 mg, 100% yield.


15

B.




Analogous to the procedure described in Example 8 Part D, Part A compound was used to prepare the title compound.

20

Example 81


The title compound was generated in analogy to the procedures described in Examples 5, 9, and 11.

5

Example 82


The title compound was generated in analogy to the procedures described in Examples 5 and 6.

10


Example 83

In analogy to the procedure described in Example 15, oxidation of Example 5 Part C compound afforded the title compound.

15

Example 84


A.



5           A mixture of Example 53 Part C compound (50 mg, 0.13 mmol),  $K_2CO_3$  (26 mg, 0.19 mmol) and DMF (3 mL) was treated with ethyl iodide (60 mg, 0.38 mmol) and the mixture stirred for 18 h. The mixture was diluted with equal amounts of ethyl acetate and water. The organic

10 fraction was washed with water, dried ( $MgSO_4$ ) and concentrated. The remainder was purified by flash column chromatography with 20% ethyl acetate in hexane to give 48 mg (94%) of the title compound.


15           B.



The experimental procedure as described in Example 53 Part E was followed for Suzuki coupling between the above Part A compound and 3-aminophenyl boronic acid to


20 give the title compound (yield 91%).

C.



The experimental procedure as set out in Example 53 Part F was followed employing the above Part B compound to give the title compound (yield 68%).


D.



The experimental procedure as set out in Example 53 Part G was followed employing the above Part C compound to give the title compound (yield 92%).


The molecular weight and mass spectral analysis of each of the compounds prepared in Examples 1 to 84 are set out in the following tables.

Table 1. Linker variations in the oxazole series



Ex.	X	Z	MW	MS observed (M+H) <sup>+</sup>
13	-OCH(Me)-	CO <sub>2</sub> H	461.52	462
54	-OCH <sub>2</sub> -	tetrazole	471.52	472
16	-CH=CH-	CO <sub>2</sub> H	443.51	441
17	-CH <sub>2</sub> CH <sub>2</sub> -	CO <sub>2</sub> H	445.52	446
3	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	446.51	447
4	-NHCO-	CO <sub>2</sub> H	460.49	461
18	cyclopropyl	CO <sub>2</sub> H	457.53	458
25	-N(Me)CH <sub>2</sub> -	CO <sub>2</sub> H	460.54	461
15	bond	CO <sub>2</sub> H	417.47	418
14	-CH <sub>2</sub> -	CO <sub>2</sub> H	431.5	432
2	bond	tetrazole	441.5	442
1	-CH <sub>2</sub> -	tetrazole	455.52	456


Table 2. Ring substitution



Ex.	HET	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
26	A	phenyl	phenyl	3-Me	H
27	A	phenyl	phenyl	H	4'-Me
24	A	phenyl	phenyl	3-Me	H
42	A	phenyl	phenyl	4-Cl	H
41	A	2-pyridyl	2-pyridyl	H	H
44	A	c-C <sub>6</sub> H <sub>11</sub>	c-C <sub>6</sub> H <sub>11</sub>	H	H
40	A	H	phenyl	H	H
43	A	c-C <sub>6</sub> H <sub>11</sub>	phenyl	H	H
46	B	Ph	H	H	H
55	B	4-F-Ph	4-F-Ph	H	H
56	C	4-F-Ph	4-F-Ph	H	H

Ex.	X-A	MW	MS observed (M+H) <sup>+</sup>
26	-NHCH <sub>2</sub> CO <sub>2</sub> H	460.54	461
27	-NHCH <sub>2</sub> CO <sub>2</sub> H	460.54	461
24	-OCH <sub>2</sub> CO <sub>2</sub> H	461.52	462
42	-NHCH <sub>2</sub> CO <sub>2</sub> H	480.96	481
41	-OCH <sub>2</sub> CO <sub>2</sub> H	449.47	449 (M) <sup>+</sup>
44	-OCH <sub>2</sub> CO <sub>2</sub> H	459.59	460
40	-OCH <sub>2</sub> CO <sub>2</sub> H	371.4	372
43	-OCH <sub>2</sub> CO <sub>2</sub> H	453.54	454
46	-OCH <sub>2</sub> CO <sub>2</sub> H	370.41	371
55	-OCH <sub>2</sub> CO <sub>2</sub> H	482.49	483
56	-OCH <sub>2</sub> CO <sub>2</sub> H	510.54	511

Table 3. X-Z variation at the ortho position



5

Ex.	X	Z	MW	MS observed (M+H) <sup>+</sup>
31	-CH=CH-	CO <sub>2</sub> H	443.51	444
32	-CH <sub>2</sub> CH <sub>2</sub> -	CO <sub>2</sub> H	445.5	446
33	bond	CO <sub>2</sub> H	417.47	418
12	bond	tetrazole	441.5	442





Table 4. Imidazole based analogs

Ex.	R <sup>8</sup>	X	Z	MW	MS observed (M+H) <sup>+</sup>
36	Me	-OCH <sub>2</sub> -	CO <sub>2</sub> H	460.54	461
38	H	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	445.53	446
39	Me	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	459.55	460
6	Et	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	473.58	474
37	Propyl	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	487.61	488
45	-CH <sub>2</sub> CH <sub>2</sub> OH	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	489.58	490
9	Et	-CH=CH-	CO <sub>2</sub> H	470.58	471
47	-CH <sub>2</sub> CH(Me) <sub>2</sub>	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	501.63	502
48	Et	-NHCH <sub>2</sub> -	tetrazole	497.61	498
11	Et	-CH <sub>2</sub> CH <sub>2</sub> -	CO <sub>2</sub> H	472.59	473
7	Et	bond	tetrazole	468.57	469
8	Et	-OCH <sub>2</sub> -	CO <sub>2</sub> H	474.56	475
51	CH <sub>2</sub> CO <sub>2</sub> H	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	503.56	504
5	Et	-CH <sub>2</sub> -	tetrazole	482.59	483
10	Et	-CH <sub>2</sub> -	CO <sub>2</sub> H	458.57	459
83	Et	bond	CO <sub>2</sub> H	444.54	445
52	Et	-OCH <sub>2</sub> -	tetrazole	498.59	499
34	H	-OCH <sub>2</sub> -	CO <sub>2</sub> H	446.51	447
35	allyl	-OCH <sub>2</sub> -	CO <sub>2</sub> H	486.58	487
57	-CH <sub>2</sub> OMe	-OCH <sub>2</sub> -	CO <sub>2</sub> H	490.56	491
58	-CH <sub>2</sub> CH <sub>2</sub> F	-OCH <sub>2</sub> -	CO <sub>2</sub> H	492.55	493
81	-CH <sub>2</sub> CH <sub>2</sub> F	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	491.57	492
82	-CH <sub>2</sub> CH <sub>2</sub> F	-CH <sub>2</sub> CH <sub>2</sub> -	CO <sub>2</sub> H	490.58	491




Table 5. Furan, thiophene, and pyrrole derivatives



Ex.	W	R <sup>9</sup>	X	Z	MW	MS observed (M+H) <sup>+</sup>
23	O	H	-CH <sub>2</sub> -	tetrazole	454.54	455
21	N-Et	H	-NHCH <sub>2</sub> -	tetrazole	496.62	497
22	S	H	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	461.59	462
20	NH	H	-CH <sub>2</sub> -	tetrazole	453.55	454
19	O	H	-OCH <sub>2</sub> -	tetrazole	470.54	471
59	O	H	-OCH <sub>2</sub> -	CO <sub>2</sub> H	446.51	447
60	N-Et	H	-CH <sub>2</sub> -	tetrazole	481.61	482
61	O	Et	-OCH <sub>2</sub> -	CO <sub>2</sub> H	474.56	475
62	N-Et	H	-OCH <sub>2</sub> -	tetrazole	497.6	498
63	N-Et	H	-OCH <sub>2</sub> -	CO <sub>2</sub> H	473.58	474
64	NH	H	-OCH <sub>2</sub> -	CO <sub>2</sub> H	445.52	446
65	O	Et	-CH <sub>2</sub> -	tetrazole	482.59	483
66	O	-CH <sub>2</sub> OH	-OCH <sub>2</sub> -	CO <sub>2</sub> H	476.53	477
68	O	-CH <sub>2</sub> OEt	-OCH <sub>2</sub> -	CO <sub>2</sub> H	504.59	505
69	NH	Et	-OCH <sub>2</sub> -	CO <sub>2</sub> H	473.58	474
70	O	-CH(OH)Me	-OCH <sub>2</sub> -	CO <sub>2</sub> H	490.56	491
67	O	-CH <sub>2</sub> OMe	-OCH <sub>2</sub> -	CO <sub>2</sub> H	490.56	491
71	NH	Et	-CH <sub>2</sub> -	tetrazole	481.61	482
72	O	Et	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	473.58	474


Table 6. Pyrazole based analogs








Ex.	R <sup>9</sup>	X	Z	MW	MS observed (M+H) <sup>+</sup>
73	Et	-OCH <sub>2</sub> -	CO <sub>2</sub> H	474.56	475
74	Et	-CH <sub>2</sub> -	tetrazole	482.59	483
84	OEt	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	489.58	490
75	Et	-NHCH <sub>2</sub> -	CO <sub>2</sub> H	473.58	474
76	Me	-OCH <sub>2</sub> -	CO <sub>2</sub> H	460.54	461
77	Me	-CH <sub>2</sub> -	tetrazole	468.57	469
80	Et	-CH <sub>2</sub> CH <sub>2</sub> -	CO <sub>2</sub> H	472.59	473
79	Et	-CH=CH-	CO <sub>2</sub> H	470.58	471


5

Table 7. Various heterocyclic based analogs




Ex.	Het	X-Z	MW	MS observed (M+H) <sup>+</sup>
49		-OCH <sub>2</sub> CO <sub>2</sub> H	452.51	475 (M+Na) <sup>+</sup>
50		-OCH <sub>2</sub> CO <sub>2</sub> H	452.51	475 (M+Na) <sup>+</sup>

28		-OCH <sub>2</sub> CO <sub>2</sub> H	476.54	477
30		-OCH <sub>2</sub> CO <sub>2</sub> H	463.5	464
29		-NHCH <sub>2</sub> CO <sub>2</sub> H	462.51	463
78		-OCH <sub>2</sub> CO <sub>2</sub> H	475.51	476
53		-NHCH <sub>2</sub> CO <sub>2</sub> H	475.55	476


Example 85

A.




- 5           A mixture of 4-methoxy-benzonitrile (1.33 g, 10.0 mmol) and THF at  $-78^{\circ}\text{C}$  was treated with benzylmagnesium chloride in THF (5.1 mL, 10.2 mmol) over 5 min. The mixture was warmed to  $45^{\circ}\text{C}$  for 2 h and cooled to ice bath temperature. The mixture was then treated slowly with 5
- 10   mL of 3N HCl and stirred for 2 h. Extraction with ethyl acetate (3X30 mL) drying ( $\text{MgSO}_4$ ) and concentrating gave a yellow oil. The oil was purified by flash column chromatography on silica gel with 4:6
- 15   hexane:dichloromethane to give 1.54 g (68%) of the title compound as a light yellow solid.

B.




- 20           Analogous to the procedure described in Example 73 Part A, Part A compound and potassium acetate were used to prepare the title compound in 82% yield.

C.




To a stirred solution of Part B compound (1.00 g, 2.53 mmol) in N-methylpyrrolidone (NMP, 2 mL), 4-dimethylamino-pyridine (DMAP, 0.31 g, 2.53 mmol), and N,N'-diisopropylethylamine (0.89 mL, 5.10 mmol) was added freshly distilled propionic anhydride (0.64 mL, 5.10 mmol) and the reaction heated to 90°C for 2 h. The mixture was then treated every 24 h for 72 h with three additional amounts of DMAP (0.31 g, 2.53 mmol), N,N'-diisopropylethylamine (0.89 mL, 5.10 mmol) and propionic anhydride (0.64 mL, 5.10 mmol). After the final addition the mixture was stirred overnight at 90°C and cooled to room temperature. The mixture was diluted with equal portions (50 mL) of 4N HCl and ethyl acetate and equilibrated. The organic fraction was washed with brine, saturated  $\text{NHCO}_3$  and brine, dried ( $\text{MgSO}_4$ ) and concentrated to give 1.49 g of the title compound as a crude brown oil.

D.



Analogous to the procedure described in Example 73 Part C, Part C compound was used to prepare the title compound in 50% yield.


5 E.



A solution of Part D compound (1.00 g, 2.3 mmol) and tris(dibenzylidene acetone)dipalladium(0) (0.13 g, 0.14 mmol) in a mixture of 1:3:1 THF:DME:water (10 mL) was treated with 3-hydroxyphenylboronic acid (0.45 g, 3.24 mmol) followed by the addition of aqueous  $\text{Na}_2\text{CO}_3$  (6.4 mL, 1.5 M, 9.63 mmol). The mixture was heated to 50°C for 2 h, cooled to room temperature and acidified to pH=3 with 2N HCl. The mixture was diluted with equal portions of ethyl acetate and brine (50 mL) and equilibrated. The organic fraction was washed with brine and 1N HCl, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to a solid. The solid was recrystallized from hot ethyl acetate to give 700 mg (68%) of the title compound.


20

F.




A slurry of Part E compound (0.70 g, 1.57 mmol) in DMF (6 mL) was treated with t-butyl bromoacetate (0.25 mL, 1.72 mmol) and cesium carbonate (0.51 g, 1.57 mmol) and stirred for 18 h. The reaction mixture was  
5 partitioned between ethyl acetate and water. The organic layer was washed with water, and brine, dried over  $\text{MgSO}_4$  and concentrated. The crude product was purified by flash column chromatography on silica gel with 3:7 ethyl acetate/hexane to give the title compound as a white  
10 solid (0.70 g, 80%).

G.




To a solution of Part F compound (0.33 g, 0.6  
15 mmol) in dry dichloromethane (2 mL) at 0°C was added boron tribromide in dichloromethane (1M, 2.7 mL, 2.7 mmol). The solution was stirred at 0°C for 3 h and at room temperature overnight. The mixture was diluted with water followed by 1N HCl and extracted with ethyl  
20 acetate. The combined organic extracts were washed with brine, dried over  $\text{MgSO}_4$  and concentrated. The crude product was purified by preparative reverse phase chromatography to give the title compound as a white  
solid (0.05 g, 15%) and Example 86 part A compound (0.18  
25 g, 55%). MS (title compound)  $[M+H]=505$ .

### Example 86




A.



5

Following the procedure in Example 85 Part F afforded the title compound (yield 55%).

B.




10

Following the procedure as in Example 28 Part G, alkylation of Part A compound with ethyl bromoacetate and cesium carbonate afforded the title compound (15%).




C.



Following the experimental procedure as in Example  
28 Part H, hydrolysis of Part B compound afforded the  
5 title compound as a white solid (85%). MS [M+H]=491.

What is Claimed is:

1. A compound having the structure






- wherein  $R^1$  and  $R^2$  are the same or different and are  
 5 independently selected from hydrogen, alkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heteroarylalkyl, aralkyl, cycloheteroalkyl or cycloheteroalkylalkyl;  
 $R^3$  is selected from hydrogen, halogen, alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkylalkyl,  
 10 cycloalkenyl, alkylcarbonyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkenylalkyl, haloalkyl, polyhaloalkyl, cyano, nitro, hydroxy, amino, alkanoyl, alkylthio, alkylsulfonyl, alkoxycarbonyl, alkylaminocarbonyl, alkylcarbonylamino, alkylcarbonyloxy,  
 15 alkylaminosulfonyl, alkylamino, dialkylamino, all optionally substituted through available carbon atoms with 1, 2, 3, 4 or 5 groups selected from hydrogen, halo, alkyl, polyhaloalkyl, alkoxy, haloalkoxy, polyhaloalkoxy, alkoxycarbonyl, alkenyl, alkynyl, cycloalkyl,  
 20 cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, hydroxy, hydroxyalkyl, nitro, cyano, amino, substituted amino, alkylamino, dialkylamino, thiol, alkylthio, alkylcarbonyl, acyl, alkoxycarbonyl, aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl,  
 25 alkenylaminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, alkoxycarbonylamino, alkylsulfonyl, aminosulfinyl, aminosulfonyl, alkylsulfinyl, sulfonamido or sulfonyl;  
 $R^4$  is selected from hydrogen, halogen, alkyl,  
 30 alkenyl, alkynyl, alkoxy, aryl, heteroaryl, arylalkyl, heteroarylalkyl, arylalkenyl, arylalkynyl, cycloalkyl, cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl, cycloalkenyl, cycloalkynyl, alkylcarbonyl, arylcarbonyl,

cycloheteroalkyl, cycloheteroalkylalkyl,  
 cycloalkenylalkyl, polycycloalkenyl,  
 polycycloalkenylalkyl, polycycloalkynyl,  
 polycycloalkynylalkyl, haloalkyl, polyhaloalkyl, cyano,  
 5 nitro, hydroxy, amino, alkanoyl, aroyl, alkylsulfonyl,  
 arylsulfonyl, alkoxycarbonyl, aryloxy carbonyl,  
 alkylaminocarbonyl, arylaminocarbonyl,  
 alkylcarbonylamino, alkoxycarbonyloxy,  
 alkylaminosulfonyl, arylaminosulfonyl, alkylamino,  
 10 dialkylamino, all optionally substituted through  
 available carbon atoms with 1, 2, 3, 4 or 5 groups  
 selected from hydrogen, halo, alkyl, haloalkyl,  
 polyhaloalkyl, alkoxy, haloalkoxy, polyhaloalkoxy,  
 alkoxycarbonyl, alkenyl, alkynyl, cycloalkyl,  
 15 cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl,  
 aryl, heteroaryl, arylalkyl, arylcycloalkyl, arylalkenyl,  
 arylalkynyl, aryloxy, aryloxyalkyl, arylalkoxy, arylazo,  
 heteroaryloxy, heteroarylalkyl, heteroarylalkenyl,  
 heteroaryloxy, hydroxy, hydroxyalkyl, nitro, cyano,  
 20 amino, substituted amino, alkylamino, dialkylamino,  
 thiol, alkylthio, arylthio, heteroarylthio,  
 arylthioalkyl, alkylcarbonyl, arylcarbonyl, acyl,  
 arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl,  
 alkynylaminocarbonyl, alkylaminocarbonyl,  
 25 alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy,  
 alkylcarbonylamino, arylcarbonylamino,  
 alkoxycarbonylamino, arylsulfinyl, arylsulfinylalkyl,  
 arylsulfonyl, alkylsulfonyl, aminosulfinyl,  
 aminosulfonyl, arylsulfonylamino,  
 30 heteroarylcarbonylamino, heteroarylsulfinyl,  
 heteroarylthio, heteroarylsulfonyl, alkylsulfinyl,  
 sulfonamido or sulfonyl;

X is a bond or a linker group selected from  $(CH_2)_n$ ,  
 $O(CH_2)_n$ ,  $S(CH_2)_n$ , cycloalkylene,  $N(R^5)(CH_2)_n$ ,  $NHCO$ , or  
 35  $CH=CH$  where  $n = 0-5$  and  $R^5$  is hydrogen, alkyl, or  
 alkanoyl;

Z is CO<sub>2</sub>H or tetrazole of the formula  
its tautomer; and






the group  represents a heteroaryl group or cycloheteroalkyl group which may further be optionally substituted with one or two groups, which may be the same or different and are independently selected from alkyl, alkenyl, hydroxyalkyl, keto, carboxyalkyl, carboxy, cycloalkyl, alkoxy, formyl, alkanoyl, alkoxyalkyl or alkoxycarbonyl,

- including all stereoisomers thereof;  
and a pharmaceutically acceptable salt thereof, or a prodrug ester thereof;

with the provisos that


- (1)  $n \neq 0$  when Z is CO<sub>2</sub>H and X is O(CH<sub>2</sub>)<sub>n</sub>, S(CH<sub>2</sub>)<sub>n</sub> or  
N(R<sup>5</sup>)(CH<sub>2</sub>)<sub>n</sub>; and



(2) when  is , then X-Z may not be O-lower alkylene-CO<sub>2</sub>H or -O-lower alkylene-CO<sub>2</sub>alkyl when R<sup>1</sup> and R<sup>2</sup> are both aryl or substituted aryl and R<sup>3</sup> and R<sup>4</sup> are each hydrogen.


2. The compound as defined in Claim 1 wherein R<sup>3</sup> and R<sup>4</sup> are the same or different and are independently selected from hydrogen, halogen, alkyl, alkoxy, alkylthio, haloalkyl, CF<sub>3</sub>, cyano, hydroxy, or nitro.

3. The compound as defined in Claim 1 wherein




includes 1 to 3 heteroatoms.

4. The compound as defined in Claim 1 wherein




is a 5-membered heteroaryl group or a 5-membered cycloheteroalkyl group.

5. The compound as defined in Claim 1 wherein




5 is a heteroaryl group.

6. The compound as defined in Claim 1 wherein



is




10









where  $R^8$  is selected from H, alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, alkenyl, and

$R^9$  is selected from H, alkyl, alkenyl, formyl,   
 15  $CO_2$ (lower alkyl), hydroxyalkyl, alkoxyalkyl,  $CO$ (alkyl), carboxyalkyl, haloalkyl, alkenyl or cycloalkyl.

7. The compound as defined in Claim 6 wherein



is


















or

8. The compound as defined in Claim 1 wherein  $R^1$  and  $R^2$  are the same or different and are independently selected from aryl, cycloalkyl, heteroaryl or hydrogen.


9. The compound as defined in Claim 1 wherein  $R^1$  and  $R^2$  are the same or different and are independently selected from phenyl, cyclohexyl, hydrogen or pyrido.

10. The compound as defined in Claim 1 wherein  $R^3$  and  $R^4$  are the same or different and are independently selected from hydrogen, alkyl or halogen.



11. The compound as defined in Claim 1 wherein -X-  
Z is



or



12. The compound as defined in Claim 1 wherein



5 where  $R^8$  is H, lower alkyl, fluoroalkyl, or alkoxyalkyl, and  $R^9$  is H, lower alkyl, fluoroalkyl, alkoxy or hydroxyalkyl;



$R^1$  and  $R^2$  are the same or different and are independently selected from phenyl or substituted phenyl;

10  $R^3$  and  $R^4$  are the same or different are independently selected from H, halo, alkyl or alkoxy;

X is  $\text{OCH}_2$ ,  $\text{NHCH}_2$ ,  $\text{CH}_2$  or  $\text{CH}_2\text{CH}_2$ ; and

Z is  $\text{CO}_2\text{H}$  or tetrazole.



13. The compound as defined in Claim 1 where



20 where  $R^8$  is H, lower alkyl or fluoroalkyl, and  $R^9$  is H, lower alkyl, fluoroalkyl, or alkoxy;

$R^1$  and  $R^2$  are each phenyl;  
 $R^3$  and  $R^4$  are each H;  
X is  $OCH_2$ ,  $CH_2$  or  $NHCH_2$ ; and  
Z is  $CO_2H$  or tetrazole.


5 14. The compound as defined in Claim 1 wherein




$R^3$  is H

$R^4$  is H

10 and -X-Z is






15. The compound as defined in Claim 1 which is










15








5















5









10















5











10
















10






5







16. The compound as defined in Claim 1 which is



17. A pharmaceutical composition comprising a  
 10 compound as defined in Claim 1 and a pharmaceutically  
 acceptable carrier therefor.

18. A pharmaceutical combination comprising an aP2  
 inhibitor compound as defined in Claim 1 and an  
 antidiabetic agent other than an aP2 inhibitor, an anti-  
 obesity agent, a lipid-lowering agent, an anti-  
 15 hypertensive agent, an anti-platelet agent and/or an  
 anti-infective agent.

19. The pharmaceutical combination as defined in Claim 18 comprising said  $\alpha$ P2 inhibitor compound and an antidiabetic agent.

20. The combination as defined in Claim 19 wherein  
5 the antidiabetic agent is 1, 2, 3 or more of a biguanide, a sulfonyl urea, a glucosidase inhibitor, a PPAR  $\gamma$  agonist, a PPAR  $\alpha/\gamma$  dual agonist, an SGLT2 inhibitor, a DP4 inhibitor, an insulin sensitizer, a glucagon-like peptide-1 (GLP-1), insulin and/or a meglitinide.

10 21. The combination as defined in Claim 20 wherein the antidiabetic agent is 1, 2, 3 or more of metformin, glyburide, glimepiride, glipyrizide, glipizide, chlorpropamide, gliclazide, acarbose, miglitol, pioglitazone, troglitazone, rosiglitazone, insulin, Gl-  
15 262570, isaglitazone, JTT-501, NN-2344, L895645, YM-440, R-119702, AJ9677, repaglinide, nateglinide, KAD1129, AR-HO39242, GW-409544, KRP297, AC2993, LY315902, and/or NVP-DPP-728A.

22. The combination as defined in Claim 19 wherein  
20 the compound is present in a weight ratio to the antidiabetic agent within the range from about 0.01 to about 100:1.

23. The combination as defined in Claim 18 wherein the anti-obesity agent is a beta 3 adrenergic agonist, a  
25 lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta compound, and/or an anorectic agent.

24. The combination as defined in Claim 23 wherein the anti-obesity agent is orlistat, ATL-962, AJ9677,  
30 L750355, CP331648, sibutramine, topiramate, axokine, dexamphetamine, phentermine, phenylpropanolamine, and/or mazindol.

25. The combination as defined in Claim 18 wherein the lipid lowering agent is an MTP inhibitor, an HMG CoA  
35 reductase inhibitor, a squalene synthetase inhibitor, a fibric acid derivative, an upregulator of LDL receptor activity, a lipoxygenase inhibitor, or an ACAT inhibitor.

26. The combination as defined in Claim 25 wherein the lipid lowering agent is pravastatin, lovastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, nisvastatin, visastatin, fenofibrate, gemfibrozil,  
5 clofibrate, avasimibe, TS-962, MD-700, and/or LY295427.

27. The combination as defined in Claim 25 wherein the aP2 inhibitor is present in a weight ratio to the lipid-lowering agent within the range from about 0.01 to about 100:1.

10 28. The combination as defined in Claim 18 wherein the anti-hypertensive agent is an ACE inhibitor, a vasopeptidase inhibitor, an angiotensin-II antagonist, a calcium-channel blocker, an alpha-blocker, a beta-blocker, a potassium channel opener, a centrally acting  
15 alpha agonist, and/or a diuretic.

29. The combination as defined in Claim 28 wherein the anti-hypertensive agent is omapatrilat, [S-(R\*,R\*)]-hexahydro-6-[(2-mercapto-1-oxo-3-phenylpropyl)amino]-2,2-dimethyl-7-oxo-1H-azepine-1-acetic acid, lisinopril,  
20 enalapril, quinapril, benazepril, fosinopril, ramipril, captopril, enalaprilat, moexipril, trandolapril, perindopril, losartan, valsartan, irbesartan, candesartan, telmisartan, amlodipine, diltiazem, nifedipine, verapamil, felodipine, nisoldipine,  
25 isradipine, nicardipine, terazosin, doxazosin, prazosin, nadolol, propranolol, metoprolol, atenolol, carvedilol, sotalol, hydrochlorthiazide, torasemide, furosemide, spironolactone, indapamide, clonidine and/or guanfacine.

30 30. The combination as defined in Claim 18 wherein the anti-platelet agent is aspirin, clopidogrel, ticlopidine, abciximab, tirofiban, eptifibatide, anagrelide and/or dipyridamole.

31. The combination as defined in Claim 18 wherein the anti-infective is azithromycin, gatifloxacin,  
35 ciprofloxacin, levofloxacin, or trovafloxacin.

32. A method for treating insulin resistance, hyperglycemia, hyperinsulinemia, or elevated blood levels of free fatty acids or glycerol, obesity, hypertriglyceridemia, Syndrome X, diabetic complications,  
5 or atherosclerosis which comprises administering to a mammalian species in need of treatment a therapeutically effective amount of a pharmaceutical composition as defined in Claim 17.

33. A method for treating Crohn's disease,  
10 ulcerative colitis, rheumatoid arthritis, chronic obstructive pulmonary disease, emphysema or systemic lupus erythematosus, which comprises administering to a human patient in need of treatment a therapeutically effective amount of a compound as defined in Claim 1.

15 34. A method for treating Crohn's disease, ulcerative colitis, rheumatoid arthritis, chronic obstructive pulmonary disease, emphysema, or systemic lupus erythematosus, which comprises administering to a human patient in need of treatment a therapeutically  
20 effective amount of a compound which inhibits  $\alpha 2$ .

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/07417

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)


U.S. : 548/235, 236, 254, 264.6, 335.1, 341.5, 342.5; 514/374, 381, 384, 399, 400

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
CHEMICAL ABSTRACTS. INDEX CHemicus; CURRENT ABSTRACTS OF CHEMISTRYElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
NONE**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,756,527 A (MJALLI et al.) 26 May 1998 (26.05.1998), see entire document.	1-34
A	US 5,811,445 A (CORBIER et al.) 22 September 1998 (22.09.1998), see entire document.	1-34
A, E	US 6,080,870 (ANTHONY et al.) 27 June 2000 (27.06.2000), see entire document.	1-34

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 14 JULY 2000	Date of mailing of the international search report <b>14 AUG 2000</b>
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer FLOYD D. HIGEL  Telephone No. (703) 308-1235

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/07417

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

A61K 31/41, 31/4164, 31/4178, 31/4196, 31/4245; C07D 233/54, 233/56, 233/60, 233/68; 249/12; 254/10; 263/30, 263/34; 403/00

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

548/235, 236, 254, 264.6, 335.1, 341.5, 342.5; 514/374, 381, 384, 399, 400